

The joint spectral radius for semigroups generated by switched differential algebraic equations

Stephan Trenn* and **Fabian Wirth****

* Technomathematics group, University of Kaiserslautern, Germany

** Department for Mathematics, University of Würzburg, Germany

SIAM Conference on Applied Linear Algebra
Valencia, Spain, 18.06.2012



Content



- 1 Introduction
- 2 Evolution operator and its semigroup
- 3 Converse Lyapunov theorem and Barabanov norm
- 4 Conclusions

Switched DAEs



Linear switched DAE (differential algebraic equation)

(swDAE)

$$E_{\sigma(t)}\dot{x}(t) = A_{\sigma(t)}x(t)$$

or short

$$E_{\sigma}\dot{x} = A_{\sigma}x$$

with

- switching signal $\sigma : \mathbb{R} \rightarrow \{1, 2, \dots, P\}$
 - piecewise constant, right-continuous
 - locally finitely many jumps (no Zeno behavior)
- matrix pairs $(E_1, A_1), \dots, (E_P, A_P)$
 - $E_p, A_p \in \mathbb{R}^{n \times n}$, $p = 1, \dots, P$
 - (E_p, A_p) regular, i.e. $\det(E_p s - A_p) \not\equiv 0$
 - impulse-free solutions (but jumps are allowed!)

Question

Growth rate and extremal norms for $E_{\sigma}\dot{x} = A_{\sigma}x \forall \sigma$



Solution formula



Theorem (A^{diff} and Π , Tanwani & T. 2010)

Let (E, A) be regular and consider

$$E\dot{x} = Ax \quad \text{on } [0, \infty)$$

$\Rightarrow \exists$ unique *consistency projector* Π and unique *flow matrix* A^{diff} :

$$x(0) = \Pi x(0-)$$

$$\dot{x} = A^{\text{diff}} x \quad \text{on } (0, \infty)$$

Furthermore, $A^{\text{diff}} \Pi = \Pi A^{\text{diff}}$.

Corollary (Solution formula for switched DAE)

Any solution of the switched DAE $E_\sigma \dot{x} = A_\sigma x$ has the form

$$x(t) = e^{A_k^{\text{diff}}(t-t_k)} \Pi_k e^{A_{k-1}^{\text{diff}}(t_k-t_{k-1})} \Pi_{k-1} \dots e^{A_1^{\text{diff}}(t_2-t_1)} \Pi_1 e^{A_0^{\text{diff}}(t_1-t_0)} \Pi_0 x(t_0-)$$

Switched ODEs with jumps



Corollary

x solves $E_\sigma \dot{x} = A_\sigma x$ on $[0, \infty) \Leftrightarrow x$ solves *switched ODE with jumps*

$$\begin{aligned} \dot{x} &= A_{p_i}^{\text{diff}} x \text{ on } [t_i, t_{i+1}) \\ x(t_i) &= \Pi_{p_i} x(t_i-), \quad i \in \mathbb{N} \end{aligned}$$

where $0 = t_0, t_1, \dots$, are the switching times of σ and $\sigma|_{[t_i, t_{i+1})} \equiv p_i$

Impulse freeness assumption

Above solution characterization only valid when switched DAE produces no Dirac impulses in x .

Theorem (Impulse freeness characterization, T. 2009)

$E_\sigma \dot{x} = A_\sigma x$ has only impulse free solutions $\forall \sigma \Leftrightarrow$

$$\forall p, q \in \{1, \dots, P\} : E_q (I - \Pi_q) \Pi_p = 0$$

Evolution operator



Consider in the following **switched ODE with jumps**

$$\begin{aligned} \dot{x} &= A_i x \text{ on } [t_i, t_{i+1}) \\ x(t_i) &= \Pi_i x(t_i-), \quad i \in \mathbb{N} \end{aligned}$$

where $0 = t_0 < t_1 < t_2 < \dots$ and

$$(A_i, \Pi_i) \in \mathcal{M} \subseteq \{ (A, \Pi) \mid A\Pi = \Pi A, \Pi = \Pi^2 \} \text{ compact}$$

Solutions:

$$x(t) = e^{A_k(t-t_k)} \Pi_k e^{A_{k-1}(t_k-t_{k-1})} \Pi_{k-1} \dots e^{A_1(t_2-t_1)} \Pi_1 e^{A_0(t_1-t_0)} \Pi_0 x(t_0-)$$

Definition (Set of all evolutions with fixed time span $t \geq 0$)

$$\mathcal{S}_t := \left\{ \prod_{i=0}^k e^{A_i \tau_i} \Pi_i \mid (A_i, \Pi_i) \in \mathcal{M}, \sum_{i=0}^k \tau_i = t, \tau_i > 0, \tau_k \geq 0 \right\}$$

Semi group property



Lemma (Semi group)

The set

$$\mathcal{S} := \bigcup_{t>0} \mathcal{S}_t$$

is a semi group with

$$\mathcal{S}_{s+t} = \mathcal{S}_s \mathcal{S}_t := \{ \Phi_s \Phi_t \mid \Phi_s \in \mathcal{S}_s, \Phi_t \in \mathcal{S}_t \}$$

Need **commutativity** to show “ \subseteq ”:

$$e^{A\tau} \Pi = e^{A(\tau-\tau')} \overset{\curvearrowright}{e^{A\tau'}} \Pi \Pi = e^{A(\tau-\tau')} \Pi e^{A\tau'} \Pi$$

for any $(A, \Pi) \in \mathcal{M}$ and $0 < \tau' < \tau$

Exponential growth bound



Definition (Exponential growth bound)

For $t > 0$ the *exponential growth bound* of $E_\sigma \dot{x} = A_\sigma x$ is

$$\lambda_t(\mathcal{S}_t) := \sup_{\Phi_t \in \mathcal{S}_t} \frac{\ln \|\Phi_t\|}{t} \in \mathbb{R} \cup \{-\infty, \infty\}$$

Definition implies for all solutions x of $E_\sigma \dot{x} = A_\sigma x$:

$$\|x(t)\| = \|\Phi_t x(0-)\| \leq \|\Phi_t\| \|x(0-)\| \leq e^{\lambda_t(\mathcal{S}_t)t} \|x(0-)\|$$

Difference to switched ODEs without jumps

$\lambda_t(\mathcal{S}_t) = \pm\infty$ is possible!




All jumps are trivial, i.e. $\Pi_p = 0 \Rightarrow \lambda_t(\mathcal{S}_t) = -\infty$

Infinite exponential growth bound



Example:

$$(E_1, A_1) = \left(\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 0 & -1 \end{bmatrix} \right) \quad (E_2, A_2) = \left(\begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right)$$



For small dwell times: $\Phi_t \approx (\Pi_1 \Pi_2)^k = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}^k = 2^{k-1} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

Existence of exponential growth rate



Theorem (Boundedness of \mathcal{S}_t)

\mathcal{S}_t is *bounded* \Leftrightarrow the set of jump *projectors* is *product bounded*

Reminder:

$$\mathcal{S}_t := \left\{ \prod_{i=0}^k e^{A_i \tau_i} \Pi_i \mid (A_i, \Pi_i) \in \mathcal{M}, \sum_{i=0}^k \tau_i = \Delta t, \tau_i > 0, \tau_k \geq 0 \right\}$$

Theorem (Exponential growth rate well defined)

Let the jump projectors be *product bounded* and not all be trivial, then the (*upper*) *Lyapunov exponent*

$$\lambda(\mathcal{S}) := \lim_{t \rightarrow \infty} \lambda_t(\mathcal{S}_t) = \lim_{t \rightarrow \infty} \sup_{\Phi_t \in \mathcal{S}_t} \frac{\|\Phi_t\|}{t}$$

of the semi-group \mathcal{S} is *well defined* and *finite*.

Connection to the generalized spectral radius



Observation: x solves switched ODE \Leftrightarrow

$$x(t+1) \in \{ \Phi x(t) \mid \Phi \in \mathcal{S}_1 \}$$

Definition (Generalized spectral radius)

For $k \in \mathbb{N}$ define the *discrete growth rate*

$$\rho_k(\mathcal{S}_1) := \sup_{\Phi_i \in \mathcal{S}_1} \|\Phi_k \Phi_{k-1} \cdots \Phi_1\|^{1/k}.$$

The **generalized spectral radius** is

$$\rho(\mathcal{S}_1) := \lim_{k \rightarrow \infty} \rho_k(\mathcal{S}_1).$$

Clearly, $\ln \rho_k(\mathcal{S}_1) = \sup_{\Phi \in \mathcal{S}_k} \frac{\ln \|\Phi\|}{k} = \lambda_k(\mathcal{S}_k)$ and therefore

$$\lambda(\mathcal{S}) = \ln \rho(\mathcal{S}_1)$$

Contents



- 1 Introduction
- 2 Evolution operator and its semigroup
- 3 Converse Lyapunov theorem and Barabanov norm**
- 4 Conclusions



Converse Lyapunov theorem for switched DAEs

Consider again

$$E_\sigma \dot{x} = A_\sigma x \quad (\text{swDAE})$$

with corresponding semigroup \mathcal{S}_t .

(swDAE) uniformly exponentially stable

$$:\Leftrightarrow \exists M \geq 1, \mu > 0 : \|x(t)\| \leq M e^{-\mu t} \|x(0-)\| \quad \forall t \geq 0$$

$$\Rightarrow \lambda(\mathcal{S}) \leq -\mu < 0.$$

Definition (Lyapunov norm)

For $\varepsilon > 0$ define

$$\| \|x\| \|_\varepsilon := \sup_{t>0} \sup_{\Phi_t \in \mathcal{S}_t} e^{-(\lambda(\mathcal{S})+\varepsilon)t} \|\Phi_t x\|$$

Theorem (Converse Lyapunov theorem, T. & Wirth 2012)

(swDAE) is uniformly exponentially stable $\forall \sigma$

$\Rightarrow V = \| \cdot \|_\varepsilon$ is **Lyapunov function** for sufficiently small $\varepsilon > 0$

In particular: $V(\Pi x) \leq V(x)$ for all projectors Π

Barabanov norm



Definition (Barabanov norm)

$\|\cdot\|$ is called **Barabanov norm** for \mathcal{S} , iff

- ① $\|\Phi_t x^0\| \leq e^{\lambda t} \|x^0\|, \quad \Phi_t \in \mathcal{S}_t$
- ② $\forall x^0 \in \mathbb{R}^n \exists \bar{\Phi}_t \in \bar{\mathcal{S}}_t: \|\bar{\Phi}_t x^0\| = e^{\lambda t} \|x^0\|$

In particular, every Barabanov norm with $\lambda < 0$ defines a Lyapunov function

Theorem (Existence of Barabanov norm)

Assume \mathcal{S} is **irreducible**, i.e. $\mathcal{S}\mathcal{M} \subseteq \mathcal{M}$ implies $\mathcal{M} = \emptyset$ or $\mathcal{M} = \mathbb{R}^n$.

Then the following are **equivalent**:

- ① The consistency projectors are **product bounded**
- ② The **Lyapunov exponent** $\lambda(\mathcal{S})$ is **bounded**
- ③ There **exists a Barabanov norm** with $\lambda = \lambda(\mathcal{S})$

Construction of Barabanov norm



Construction of Barabanov norm similar as in (Wirth 2002, LAA):

$$\mathcal{S}_\infty := \bigcap_{T \geq 0} \overline{\bigcup_{t \geq T} e^{-\lambda(S)t} \mathcal{S}_t}$$

is a compact nontrivial semigroup, the **limit semigroup**.

$$\|x\| := \max \{ \|Sx\| \mid S \in \mathcal{S}_\infty \}$$

is the sought Barabanov norm.

Conclusions



- Studied switched DAEs $E_\sigma \dot{x} = A_\sigma x$
- Key observation:

$$x(t) = e^{A_k^{\text{diff}}(t-t_k)} \Pi_k \dots e^{A_1^{\text{diff}}(t_2-t_1)} \Pi_1 e^{A_0^{\text{diff}}(t_1-t_0)} \Pi_0 x(t_0-)$$

- Flow set

$$\mathcal{S}_t := \left\{ \prod_{i=0}^k e^{A_i^{\text{diff}} \tau_i} \Pi_i \mid (A_i^{\text{diff}}, \Pi_i) \in \mathcal{M}, \sum_{i=0}^k \tau_i = \Delta t, \tau_i > 0 \right\}$$

- **Product boundedness** of consistency projectors necessary and sufficient for boundedness of \mathcal{S}_t
- **Converse Lyapunov theorem**
- Construction of **Barabanov norm** in irreducible case