

## Regularity of distributional differential algebraic equations

Stephan Trenn

August 6, 2009

**Abstract** Time-varying differential algebraic equations (DAEs) of the form  $E\dot{x} = Ax + f$  are considered. The solutions  $x$  and the inhomogeneities  $f$  are assumed to be distributions (generalized functions). As a new approach, distributional entries in the time-varying coefficient matrices  $E$  and  $A$  are allowed as well. Since a multiplication for general distributions is not possible, the smaller space of piecewise-smooth distributions is introduced. This space consists of distributions which could be written as the sum of a piecewise-smooth function and locally finite Dirac impulses and derivatives of Dirac impulses. A restriction can be defined for the space of piecewise-smooth distributions, this restriction is used to study DAEs with inconsistent initial values; basically, it is assumed that some past trajectory for  $x$  is given and the DAE is activated at some initial time. If this initial trajectory problem has a unique solution for all initial trajectories and all inhomogeneities, then the DAE is called regular. This generalizes the regularity for classical DAEs (i.e. a DAE with constant coefficients). Sufficient and necessary conditions for the regularity of distributional DAEs are given.

**Keywords** differential algebraic equations · distributional solutions · regularity

**Mathematics Subject Classification (2000)** 34A09 · 34A12 · 46F99 · 34A37

## 1 Introduction

### 1.1 Aims and main results of the paper

A differential algebraic equation (DAE) is an equation of the form

$$E\dot{x} = Ax + f$$

where  $E, A$  are in general time-varying rectangular matrices and  $f$  is some inhomogeneity. The aim of this paper is to develop a solution theory for *distributional* DAEs, i.e. distributions as introduced by Schwartz [26] are considered as solutions  $x$ , as inhomogeneities  $f$  and as entries of the coefficient matrices  $E$  and  $A$ . This framework encompasses time-varying DAEs where jumps are allowed in the coefficients. For general distributions a multiplication within the space of distributions is not possible; therefore, a smaller space, the space of *piecewise-smooth distributions*  $\mathbb{D}_{\text{pwc}^\infty}$ , is introduced (Definition 1). It is shown that it is possible to define a multiplication within  $\mathbb{D}_{\text{pwc}^\infty}$  (Theorem 4) and also a distributional restriction (Definition 8), the latter is used to formulate initial trajectory problems (Definition 15) which allows to study solutions with inconsistent initial values.

For the constant coefficient case, it is well known that *regularity* of the matrix pair  $(E, A)$  is an important concept with respect to existence and uniqueness of solutions; this concept is generalized for distributional DAEs (Definition 16). Necessary conditions for regularity (Theorem 18, Theorem 19) as well as sufficient conditions (Theorem 21, Theorem 24) are given. The sufficient conditions can be summarized in the condition that the matrix pair  $(E, A)$  can be put into a generalized Weierstraß form (Corollary 26).

### 1.2 Motivation for studying distributional DAEs

DAEs of the form  $E\dot{x} = Ax + f$  arise for example in modeling electrical circuits, mechanical and chemical systems (see e.g. [19, I.1.3]), in particular, if these models are generated automatically. If the inhomogeneity is not continuous (for example, if it is generated by a switching controller) then, even for constant coefficients, a classical solution does in general not exist because it involves the derivative of the inhomogeneity. To resolve this problem distributional solutions are considered.

The motivation to consider distributional entries in the coefficients follows from the need to study switched DAEs, which appear in case of possible structural changes in the system. For an overview on classical switched systems and for further motivation, see e.g. [21]. Inconsistent initial values can also be interpreted as a result of switching. Switching yields that the coefficient matrices  $E$  and  $A$  are time-varying and not continuous. Equivalent system description and normal forms play an important role for the analysis of DAEs. Two canonical transformations which do not change the qualitative solution behavior are, firstly, multiplication of  $E\dot{x} = Ax + f$  with some invertible (time-varying)

matrix  $S$  from the left and, secondly, a coordinate transformation  $x = Tz$  for some invertible (time-varying) matrix  $T$  with corresponding new variable  $z$ . This yields the equivalent DAE

$$SET\dot{z} = (SAT - SET')z + Sf.$$

If  $E$  and  $A$  are time-varying and not continuous (e.g. piecewise constant for switched systems), then it is reasonable to assume that  $S$  and  $T$  may also be discontinuous. Hence  $T'$  only makes sense in the distributional sense. This motivates distributional entries in the coefficient matrices. Furthermore, linear impulsive systems (see e.g. [20]) can be rewritten as a distributional ODE  $\dot{x} = Ax + f$  with distributional entries in  $A$ . For further motivation see also the switched electrical circuit example in the Appendix.

### 1.3 Results in the literature

Distributional solutions for linear DAEs were considered already in [1] and [30], mainly to deal with inconsistent initial values, but no general distributional solution theory was introduced, problems like evaluations of distributions at a certain point (which is needed to speak of initial values) were not addressed.

A first rigorous distributional solution theory was given by Cobb in [2], he introduced “piecewise continuous distributions” which encompass piecewise-smooth distributions. However, the space of piecewise continuous distributions is not closed under differentiation, and since Cobb seems to have overlooked this fact, some of the results in [2] might need a reformulation.

Another rigorous approach to study distributional solutions is based on the space of “impulsive smooth distributions” which has its origin in [13], which itself is based on [12] (but the space is not explicitly mentioned in the latter). Most papers which use the space of impulsive-smooth distribution consider only the interval  $[0, \infty)$  for solutions. This makes it conceptually difficult to formulate inconsistent initial values (because there is no past). This drawback does not exist for the space of impulsive-smooth distributions as defined in [24] (see also [25] and [19]), which are a subspace of the space of piecewise-smooth distributions with the property that jumps and impulses can only occur at  $t = 0$ .

Impulsive-smooth distributions are used in several papers as underlying solution space, e.g. for studying solvability and consistency of non-regular classical DAEs [8,9], behavioral equivalence [16] and infinite eigenvalues of the matrix polynomial  $A(s)$  with corresponding differential equation  $A(\frac{d}{dt})x = 0$  [29,28].

Based on impulsive-smooth distributions, a space of distributions is defined in [10,11] which seems to be identical with the space of piecewise-smooth distributions. This space is then used to study “multimode” (or switched) systems in the behavioral setup. Since no regularity assumption is made on the system the results are not directly comparable to the results obtained here, in particular, in [10] it is necessary to give a jump-map a priori which is in

contrast to the theory developed here, because for regular switched DAEs the jump already follows from the coefficient matrices alone. Furthermore, in [10] Dirac impulses are identified with polynomials in  $\mathbb{R}[p]$  with polynomial variable  $p$  and it is not clear how Dirac impulses at different times are distinguished.

Piecewise-smooth distributions were used as an underlying solution space for time-invariant higher order Rosenbrock systems in [14], “time-varying” topics like inconsistent initial values and switched systems were not addressed. There is no literature on DAEs with distributional coefficient matrices.

There have been numerous approaches to define a multiplication for distributions. König [17] enlarged the space of distributions to define a multiplication, Fuchssteiner [6] introduced the space of “almost bounded” distributions, see also [7]. This space is very similar, but not identical, to the space of piecewise-smooth distributions and he defined an associative multiplication which ensures that the product rule for differentiation is fulfilled. This non-commutative multiplication is identical to the multiplication defined in this paper for piecewise-smooth distributions, although the approach is very different. In [31] a commutative but non-associative multiplication was defined for another subspace of distributions. Finally, there are several textbooks on the topic of multiplications of distributions [3, 15, 23], but the results are either too general (for example results on non-associative multiplications) or too restrictive (for example results on commutative multiplications) for the purpose of this paper. In Remark 7 an additional literature review is carried out with the focus on the definition of the square of the Dirac-impulse.

One should not confuse the “multiplication” of distribution as in e.g. [13] with the multiplication as above, because the former is a convolution and is used to rewrite the derivative as a convolution with the derivative of the Dirac impulse. This view point allows for a nice algebraic analysis of time-invariant DAEs, but seems unfruitful for general time-varying DAEs.

#### 1.4 Organization of the paper and notation

The paper is organized as follows. In Section 2 the classical distribution theory is revised and the space of piecewise-smooth distributions is introduced. For piecewise-smooth distributions, the Fuchssteiner multiplication and a distributional restriction are defined. Some calculation rules are presented. In Section 3 regularity of a distributional DAE is defined, this is strongly related to solvability of a DAE and the uniqueness of solutions. Necessary and sufficient conditions for regularity are presented. Finally, in Section 5 a simple switched electrical circuit is studied to illustrate the developed distributional solution theory.

To improve readability all proofs are carried out in the Appendix.

The following notation is used throughout the paper.  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{R}$  are the natural numbers, integers and real number, respectively. The space of functions  $\alpha$  :

$\mathbb{R} \rightarrow \mathbb{R}$  which are smooth (i.e. arbitrarily often differentiable) is denoted by  $\mathcal{C}^\infty$ . The restriction  $f_M : \mathbb{R} \rightarrow \mathbb{R}$  of some function  $f : \mathbb{R} \rightarrow \mathbb{R}$  on some set  $M \subseteq \mathbb{R}$  is defined by  $f_M := \mathbb{1}_M f$ , where  $\mathbb{1}_M : \mathbb{R} \rightarrow \{0, 1\}$  is the indicator function of  $M$  (i.e.  $\mathbb{1}_M(t) = 1$  if, and only if,  $t \in M$ ), in particular, the restricted function  $f_M$  is still defined on the whole of  $\mathbb{R}$ . A function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is called locally integrable if, and only if,  $f$  is (Lebesgue-)measurable and for every compact set  $K \subseteq \mathbb{R}$  the (Lebesgue-)integral  $\int_K |f|$  is finite. The space of distributions is  $\mathbb{D}$  and the space of piecewise-smooth distribution is  $\mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  (see the later definitions).

## 2 Distributions

### 2.1 Review of classical distribution theory

Basis knowledge of distribution theory as introduced by Schwartz [26] is assumed and is only summarized without proofs in the following paragraph.

The space of *test functions*  $\mathcal{C}_0^\infty \subseteq \mathcal{C}^\infty$  is the set of all functions  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$  which are *smooth* and which have *bounded support* (the support  $\text{supp } \varphi \subseteq \mathbb{R}$  of  $\varphi$  is the closure of  $\{t \in \mathbb{R} \mid \varphi(t) \neq 0\}$ ). The space  $\mathcal{C}_0^\infty$  is a topological vector space, and a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\mathcal{C}_0^\infty$  converges to zero (in this topology) if, and only if, there exists a compact set  $M \subseteq \mathbb{R}$  with  $\text{supp } \varphi_n \subseteq M$  for all  $n \in \mathbb{N}$  and, for each  $i \in \mathbb{N}$ , the sequence of the  $i$ -th derivatives  $(\varphi_n^{(i)})_{n \in \mathbb{N}}$  converges uniformly to zero. The space of *distributions*  $\mathbb{D}$  is the set of all *linear* and *continuous* operators  $D : \mathcal{C}_0^\infty \rightarrow \mathbb{R}$ .

The *derivative* of a distribution  $D \in \mathbb{D}$  is defined by  $D'(\varphi) := -D(\varphi')$  for all test functions  $\varphi \in \mathcal{C}_0^\infty$  and is itself a distribution. Every distribution  $D \in \mathbb{D}$  has an *antiderivative*  $H \in \mathbb{D}$ , i.e.  $H' = D$ , and all antiderivatives of  $D$  only differ by a constant distribution (the constant distribution is given by  $\varphi \mapsto c \int_{\mathbb{R}} \varphi$  for  $c \in \mathbb{R}$ ). The space of locally integrable functions  $\mathcal{L}_{1,\text{loc}}$  is injectively embedded into the space of distributions via the homomorphism

$$\mathcal{L}_{1,\text{loc}} \ni f \mapsto f_{\mathbb{D}} := \left( \varphi \mapsto \int_{\mathbb{R}} f \varphi \right) \in \mathbb{D}. \quad (1)$$

For differentiable functions  $f$ , the distributional derivative “equals” the standard derivative, i.e.  $(f')_{\mathbb{D}} = (f_{\mathbb{D}})'$ . Distributions induced by locally integrable functions via homomorphism (1) are called *regular distributions*. The well known *Dirac impulse* (also known as (Dirac-) $\delta$ -function)  $\delta_t \in \mathbb{D}$  at some time  $t \in \mathbb{R}$  is given by

$$\delta_t : \mathcal{C}_0^\infty \rightarrow \mathbb{R}, \varphi \mapsto \varphi(t)$$

and is the classical example for a distribution which is not regular.

The *support*  $\text{supp } D \subseteq \mathbb{R}$  of a distribution  $D \in \mathbb{D}$  is the complement of the largest open set on which  $D$  vanishes, i.e.

$$\text{supp } D := \mathbb{R} \setminus \bigcup \left\{ O \subseteq \mathbb{R} \mid \begin{array}{l} O \text{ open and } \forall \varphi \in \mathcal{C}_0^\infty : \\ \text{supp } \varphi \subseteq O \Rightarrow D(\varphi) = 0 \end{array} \right\}.$$

The support of the Dirac impulse  $\delta_t$  at  $t \in \mathbb{R}$  and of all its derivatives is  $\{t\}$  and, conversely, the following implication holds for all  $t \in \mathbb{R}$  and all  $D \in \mathbb{D}$

$$\text{supp } D = \{t\} \quad \Rightarrow \quad \exists n \in \mathbb{N} \exists a_0, a_1, \dots, a_n \in \mathbb{R} : D = \sum_{i=0}^n a_i \delta_t^{(i)} \quad (2)$$

and the representation is unique, i.e.  $\sum_{i=0}^n a_i \delta_t^{(i)} = \sum_{i=0}^n b_i \delta_t^{(i)}$  if, and only if,  $a_i = b_i$ ,  $i = 0, \dots, n$ . Furthermore, one can show that the only regular distribution whose support has Lebesgue measure zero (for example any countable set) is the zero distribution or in other words the support of nontrivial regular distributions is essential.

Distributions can be *multiplied by smooth functions*, i.e. for all  $\alpha \in \mathcal{C}^\infty$  the product  $\alpha D$  given by  $(\alpha D)(\varphi) := D(\alpha\varphi)$ ,  $\varphi \in \mathcal{C}_0^\infty$ , is again a distribution. It is easy to see, that the multiplication rule for the derivative holds, i.e.

$$\forall \alpha \in \mathcal{C}^\infty \quad \forall D \in \mathbb{D} : \quad (\alpha D)' = \alpha' D + \alpha D' \quad (3)$$

A simple consequence of [18, Folg. 3.24] is the following property for all  $\alpha \in \mathcal{C}^\infty$  and  $D \in \mathbb{D}$

$$\left[ \forall i \in \mathbb{N} \quad \forall t \in \text{supp } D : \alpha^{(i)}(t) = 0 \right] \quad \Rightarrow \quad \alpha D = 0. \quad (4)$$

Note that in property (4) it is not assumed that  $\text{supp } \alpha \cap \text{supp } D = \emptyset$ .

*Convergence* of a sequence of distributions is defined “point-wise”, i.e. a sequences  $(D_n)_{n \in \mathbb{N}} \in \mathbb{D}^{\mathbb{N}}$  converges to  $D \in \mathbb{D}$  if, and only if,  $D_n(\varphi) \rightarrow D(\varphi)$  for all test functions  $\varphi \in \mathcal{C}_0^\infty$ . The space  $\mathbb{D}$  is closed with respect to this convergence, i.e. if for a sequences  $(D_n)_{n \in \mathbb{N}}$  of distributions the point-wise limit exists then this limit is a distribution or, more formally,

$$\left[ \forall \varphi \in \mathcal{C}_0^\infty : \lim_{n \rightarrow \infty} D_n(\varphi) \in \mathbb{R} \right] \quad \Rightarrow \quad \lim D_n := (\varphi \mapsto \lim_{n \rightarrow \infty} D_n(\varphi)) \in \mathbb{D}. \quad (5)$$

Furthermore, for all sequences  $(D_n)_{n \in \mathbb{N}}$  of distributions,

$$\lim_{n \rightarrow \infty} D_n = D \in \mathbb{D} \quad \Rightarrow \quad \forall i \in \mathbb{N} : \lim_{n \rightarrow \infty} D_n^{(i)} = D^{(i)}. \quad (6)$$

## 2.2 Piecewise-smooth distributions

**Definition 1 (Piecewise-smooth functions and distributions)** Let the space of *piecewise-smooth functions* be given by

$$\mathcal{C}_{\text{pw}}^\infty := \left\{ \alpha = \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} \alpha_i \quad \left| \quad \begin{array}{l} \{ t_i \in \mathbb{R} \mid i \in \mathbb{Z} \} \text{ locally finite,} \\ (\alpha_i)_{i \in \mathbb{Z}} \in (\mathcal{C}^\infty)^{\mathbb{Z}} \end{array} \right. \right\}.$$

The space of piecewise-smooth distributions is defined as

$$\mathbb{D}_{\text{pw}\mathcal{C}^\infty} := \left\{ D = f_{\mathbb{D}} + \sum_{t \in T} D_t \quad \left| \quad \begin{array}{l} f \in \mathcal{C}_{\text{pw}}^\infty, \quad T \subset \mathbb{R} \text{ locally finite,} \\ \forall t \in T : D_t \in \mathbb{D} \wedge \text{supp } D_t \subseteq \{t\} \end{array} \right. \right\}$$

For a piecewise-smooth distribution  $D \in \mathbb{D}_{\text{pwC}^\infty}$  with representation  $D = f_{\mathbb{D}} + \sum_{t \in T} D_t$  the *left and right sided evaluation at  $t \in \mathbb{R}$*  is defined by

$$D(t+) := \lim_{\varepsilon \searrow 0} f(t + \varepsilon), \quad D(t-) := \lim_{\varepsilon \searrow 0} f(t - \varepsilon)$$

The *impulsive part at  $t \in \mathbb{R}$*  of the above  $D$  is defined by

$$D[t] := \begin{cases} D_t, & t \in T \\ 0, & \text{otherwise} \end{cases}$$

and the *impulsive part of  $D$*  is defined by

$$D[\cdot] := \sum_{t \in T} D[t] = \sum_{t \in T} D_t.$$

Finally,  $D_{\text{reg}} := f_{\mathbb{D}} = D - D[\cdot]$  is called the *regular part of  $D$* .

It is easy to show that the representation of piecewise-smooth distributions is unique, i.e. two piecewise-smooth distributions  $D_1, D_2 \in \mathbb{D}_{\text{pwC}^\infty}$  with corresponding representation  $D_k = f_{\mathbb{D}}^k + \sum_{t \in T^k} D_t^k$ ,  $k = 1, 2$ , are equal if, and only if,  $f^1 = f^2$ ,  $\forall t \in T^1 \cap T^2 : D_t^1 = D_t^2$ ,  $\forall t \in T^1 \setminus T^2 : D_t^1 = 0$  and  $\forall t \in T^2 \setminus T^1 : D_t^2 = 0$ . Hence the definition of left and right sided evaluation and of the impulsive part are well defined. It is important to notice the condition that the set  $T$  in the definition of  $\mathbb{D}_{\text{pwC}^\infty}$  is *locally finite*, i.e. any intersection with some compact set is finite. If this condition is not fulfilled, it is, on the one hand, not true in general that the infinite sum of distribution with point support (as in the definition of  $\mathbb{D}_{\text{pwC}^\infty}$ ) exists and, on the other hand, there are some locally infinite sums which define a distribution, but these distributions might have undesirable properties (see Remark 9).

An important motivation for introducing distributions as generalized functions is the property that every distribution has a derivative within the space of distributions. This property is preserved for the smaller space of piecewise-smooth distributions as the following proposition shows.

**Proposition 2 (Derivative of piecewise-smooth distributions)** *Let  $D \in \mathbb{D}_{\text{pwC}^\infty}$  and  $f_{\mathbb{D}} = D_{\text{reg}}$  with  $f = \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i$  for some locally finite set  $\{ t_i \in \mathbb{R} \mid i \in \mathbb{Z} \}$  and some smooth  $f_i \in \mathcal{C}^\infty$ ,  $i \in \mathbb{Z}$ . Then*

$$D' = \left( \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i' \right)_{\mathbb{D}} + \sum_{i \in \mathbb{Z}} (D(t_i+) - D(t_i-)) \delta_{t_i} + D[\cdot]'. \quad (7)$$

*In particular*

$$D \in \mathbb{D}_{\text{pwC}^\infty} \quad \Rightarrow \quad D' \in \mathbb{D}_{\text{pwC}^\infty}.$$

Piecewise-smooth distributions also have antiderivatives which are piecewise-smooth distributions again, furthermore it is possible to make the antiderivative unique.

**Proposition 3 (Unique distributional antiderivative)** For  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  and  $t_0 \in \mathbb{R}$  there exists a unique distributional antiderivative

$$H = \int_{t_0} D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$$

with  $H' = D$  and  $H(t_0-) = 0$ .

It is possible to define a multiplication for piecewise-smooth distributions as stated in the next theorem. This multiplication is a generalization of the multiplication of functions but it is not commutative anymore.

**Theorem 4 (Multiplication of piecewise-smooth distributions)** For  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  and  $t \in \mathbb{R}$  let

$$\begin{aligned} \delta_t D &:= D(t-)\delta_t, \quad \forall n \in \mathbb{N}: \delta_t^{(n+1)} D := \left( \delta_t^{(n)} D \right)' - \delta_t^{(n)} D', \\ D \delta_t &:= D(t+)\delta_t, \quad \forall n \in \mathbb{N}: D \delta_t^{(n+1)} := \left( D \delta_t^{(n)} \right)' - D' \delta_t^{(n)}. \end{aligned} \quad (8)$$

Let  $F, G \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  with representation  $F = f_{\mathbb{D}} + \sum_{t \in T_F} F[t]$  and  $G = g_{\mathbb{D}} + \sum_{t \in T_G} G[t]$  as in Definition 1. The product of  $F$  and  $G$  is defined by, using (2) and (8),

$$FG := (fg)_{\mathbb{D}} + \sum_{t \in T_F} F[t]G_{\text{reg}} + \sum_{t \in T_G} F_{\text{reg}}G[t] = (fg)_{\mathbb{D}} + F[\cdot]G_{\text{reg}} + F_{\text{reg}}G[\cdot]. \quad (9)$$

The multiplication of piecewise-smooth distributions has the following properties,  $F, G, H \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ ,  $f, g \in \mathcal{C}_{\text{pw}}^\infty$ :

- (i)  $FG \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ ,
- (ii)  $f_{\mathbb{D}}g_{\mathbb{D}} = (fg)_{\mathbb{D}}$ ,
- (iii)  $F(GH) = (FG)H$ ,  $(F+G)H = FH+GH$ ,  $F(G+H) = FG+FH$ ,
- (iv)  $(FG)' = F'G + FG'$ ,
- (v)  $\text{supp}(FG) \subseteq \text{supp} F \cap \text{supp} G$ ,

*Remark 5* Fuchssteiner [6] (see also [7]) studied the space of *almost bounded distributions*, which is very similar (but not equal) to the space of piecewise-smooth functions. He showed that the multiplication from Theorem 4 is the only one which fulfills the following conditions:

- (M1)  $\mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  is an associative differential algebra,
- (M2)  $\forall f, g \in \mathcal{C}_{\text{pw}}^\infty: (fg)_{\mathbb{D}} = f_{\mathbb{D}}g_{\mathbb{D}}$ ,
- (M3)  $\forall t \in \mathbb{R}: \mathbb{1}_{[t, \infty)_{\mathbb{D}}}\delta_t = \delta_t$ .

Hence the multiplication defined in Theorem 4 might be called *Fuchssteiner multiplication*. The motivation for introducing a multiplication for piecewise-smooth distribution was the aim to study coordinate transformations  $x = Tz$  for DAEs  $E\dot{x} = Ax$  where the time-varying matrix  $T$  has jumps. In particular  $E\dot{x} = Ax$  should be “equivalent” to  $ET\dot{z} = (A - T')z$ . To carry out this calculation (M1) must be assumed. Clearly, (M2) should hold anyway. On a first



glance, the condition (M3) might seem arbitrary. In fact, Fuchssteiner gave a description of all multiplications fulfilling (M1) and (M2). These multiplications are parametrized by a set  $M \subseteq \mathbb{R}$  and fulfill

$$\mathbb{1}_{[t, \infty)} \mathbb{D} \delta_t = \begin{cases} \delta_t, & t \in M \\ 0, & t \in \mathbb{R} \setminus M \end{cases}$$

However, only with (M3) the causality principle for DAEs of the form  $E\dot{x} = Ax$  holds. In particular, Theorem 24 shows that for any given initial trajectory  $x^0$  and initial time  $t_0 \in \mathbb{R}$  there exists a unique solution ODE  $\dot{x} = Ax$ . If one would consider another multiplication which only fulfills (M1) and (M2), but not (M3), then there would exist initial trajectory problems for distributional ODEs which are not solvable, which is, from an application-point-of-view, not desirable.

*Remark 6* From the recursive definition (8) an explicit representation can be derived easily,  $t \in \mathbb{R}$ ,  $n \in \mathbb{N}$ ,  $D \in \mathbb{D}_{\text{pwc}\infty}$ :

$$\begin{aligned} \delta_t^{(n)} D &= \sum_{i=0}^n (-1)^i \binom{n}{i} D^{(i)}(t-) \delta_t^{(n-i)}, \\ D \delta_t^{(n)} &= \sum_{i=0}^n (-1)^i \binom{n}{i} D^{(i)}(t+) \delta_t^{(n-i)}. \end{aligned} \tag{10}$$

*Remark 7 (Square of Dirac impulse)* It follows from the definition of the multiplication that  $F[\cdot]G[\cdot] = 0$  for all  $F, G \in \mathbb{D}_{\text{pwc}\infty}$  and, in particular for  $\delta := \delta_0$ ,

$$\delta^2 = 0.$$

It is interesting to compare the different approaches in the literature with respect to the square of the Dirac impulse: In [32] it is claimed that it is impossible to define this square<sup>1</sup>. A similar result is obtained in [31, Thm. 3.9], however, in the proof it is shown that the square of the Dirac impulse, if it exists, must be zero which contradicts the assumptions made in that paper. In [22] the equation  $\delta^2 - \frac{1}{\pi^2} \left(\frac{1}{x}\right)^2 = -\frac{1}{\pi^2} \frac{1}{x^2}$  is established, where the left hand side is considered as a “single entity”, this is motivated by quantum mechanics where  $\delta^2$  appears only in this context. The square of the Dirac impulse is well defined in [17], but only in a generalized space of distributions and it is shown that  $\delta^2$  is not a classical distribution. In [5] a commutative multiplication for a subspace of distributions is defined and there the square of the Dirac-impulse is zero.

The following definition introduces the restriction of piecewise-smooth distributions, which is a generalization of the restriction for functions defined by

<sup>1</sup> [32, 3.IV]: “Im besonderen ist es nicht möglich, das Quadrat der  $\delta$ -Funktion  $\delta^2$  zu bilden.”

$f_M := \mathbb{1}_M f$  for any  $f : \mathbb{R} \rightarrow \mathbb{R}$  and any  $M \subseteq \mathbb{R}$ . The distributional restriction is necessary to study *inconsistent initial values* for DAEs, because inconsistent initial values mean that the actual DAE is not valid in the past, hence it must be possible to formulate mathematically that the DAE (with distributional solutions) holds only on the time interval  $[t_0, \infty)$  for some  $t_0 \in \mathbb{R}$  (see Definition 15).

**Definition 8 (Distributional restriction)** Let  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  have the representation  $D = f_{\mathbb{D}} + \sum_{t \in T} D_t$  as in Definition 1 and let  $M \subseteq \mathbb{R}$  be a locally finite union of intervals (i.e. any compact set only intersects with finitely many intervals), then the restriction  $D_M \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  of the piecewise-smooth distribution  $D$  to the set  $M$  is

$$D_M := (f_M)_{\mathbb{D}} + \sum_{t \in T \cap M} D_t.$$

Clearly, in the above definition, the assumption that  $M \subseteq \mathbb{R}$  is the locally finite union of intervals (LFUI) ensures that  $f_M \in \mathcal{C}_{\text{pw}}^\infty$ . Furthermore,  $T \cap M$  is a locally finite set (even for arbitrary sets  $M \subseteq \mathbb{R}$ ), hence the above restriction is well defined. It is easy to see that the restriction from Definition 8 has the following properties:

(R1) The distributional restriction is a mapping

$$\{ M \subseteq \mathbb{R} \mid M \text{ is a LFUI} \} \times \mathbb{D}_{\text{pw}\mathcal{C}^\infty} \rightarrow \mathbb{D}_{\text{pw}\mathcal{C}^\infty}, \quad (M, D) \mapsto D_M$$

which is for each fixed  $M \subseteq \mathbb{R}$  a projection, i.e.  $D \mapsto D_M$  is linear and idempotent.

(R2) For  $f \in \mathcal{C}_{\text{pw}}^\infty$  and a LFUI  $M \subseteq \mathbb{R}$  the distributional restriction fulfills

$$(f_M)_{\mathbb{D}} = (f_{\mathbb{D}})_M,$$

i.e. it is a generalization of restrictions of functions.

(R3) The restriction property for trivial cases is fulfilled, i.e. for all test functions  $\varphi \in \mathcal{C}_0^\infty$ , for all distributions  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  and for all LFUIs  $M \subseteq \mathbb{R}$  the following two implications hold:

$$\begin{aligned} \text{supp } \varphi \subseteq M &\Rightarrow D_M(\varphi) = D(\varphi), \\ \text{supp } \varphi \cap M = \emptyset &\Rightarrow D_M(\varphi) = 0. \end{aligned}$$

(R4) For any pairwise disjoint family of LFUIs  $(M_i)_{i \in \mathbb{N}}$  with  $M := \bigcup_{i \in \mathbb{N}} M_i$  a LFUI and any  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  the restriction fulfills

$$D_M = \sum_{i \in \mathbb{N}} D_{M_i},$$

in particular,

$$D_{M_1 \cup M_2} = D_{M_1} + D_{M_2}.$$

Furthermore, for any disjoint LFUI sets  $M_1, M_2 \subseteq \mathbb{R}$  the restriction fulfills

$$(D_{M_1})_{M_2} = 0.$$

Note that it is crucial that only piecewise-smooth distributions are considered as the following remark shows.

*Remark 9* For general distributions it is *not possible* to define a restriction with the properties (R1)-(R4). As an example consider the following (well defined!) distribution

$$D = \sum_{n \in \mathbb{N}} d_n \delta_{d_n}, \quad d_n := \frac{(-1)^n}{n+1}, \quad n \in \mathbb{N}.$$

The restriction to the interval  $(0, \infty)$  should then be

$$D_{(0, \infty)} = \sum_{k \in \mathbb{N}} \frac{1}{2k+1} \delta_{\frac{1}{2k+1}},$$

but it is easy to see, that there exist test functions (in fact, any test function  $\varphi \in \mathcal{C}_0^\infty$  with  $\varphi(0) \neq 0$  will do) for which the infinite sum does not converge, hence the restriction is not defined.

*Remark 10* The restriction  $D_{(s,t)}$  to an open interval  $(s,t) \subseteq \mathbb{R}$  for  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  as above needs to be distinguished from the “classical” restriction to an open interval  $D|_{(s,t)}$  because the latter is only defined for test functions  $\varphi \in \mathcal{C}_0^\infty$  with  $\text{supp } \varphi \subset (s,t)$ . Remark 9 shows that for general distributions this is indeed essential. However, for piecewise-smooth distributions both restrictions (to open(!) intervals) are strongly connected as the following equality shows, for  $F, G \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ :

$$F_{(s,t)} = G_{(s,t)} \quad \Leftrightarrow \quad F|_{(s,t)} = G|_{(s,t)}.$$

Note that on the left hand side one can plug in any test function  $\varphi \in \mathcal{C}_0^\infty$  whilst on the right hand side onyl test functions  $\varphi \in \mathcal{C}_0^\infty$  with  $\text{supp } \varphi \subset (s,t)$  are allowed.

### 2.3 Calculation rules for piecewise-smooth distributions

In this subsection some calculation rules for the restriction, multiplication and differentiation of piecewise-smooth distributions are given. These will be needed in later parts of this work and are also of general interest.

**Proposition 11 (Multiplication and restriction)** *Let  $F, G \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  and  $s, t \in \mathbb{R} \cup \{\pm\infty\}$  with  $s \leq t$ , then, for any  $\varepsilon > 0$ ,*

$$\begin{aligned} (FG)_{(s,t)} &= F_{(s,t)} G_{(s,t)}, \\ (FG)_{[s,t]} &= F_{[s,t]} G_{[s,t]} + F[s] G_{(s-\varepsilon, s)}, \\ (FG)_{(s,t]} &= F_{(s,t]} G_{(s,t]} + F_{(t, t+\varepsilon)} G[t], \\ (FG)_{[s,t]} &= F_{[s,t]} G_{[s,t]} + F_{(t, t+\varepsilon)} G[t] + F[s] G_{(s-\varepsilon, s)}, \end{aligned}$$

where  $F[\pm\infty] = G[\pm\infty] = 0$ .

**Proposition 12 (Restrictions and derivatives)** For all  $-\infty \leq s \leq t \leq \infty$  and  $D \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ ,

$$\begin{aligned} (D_{[s,t]})' &= (D')_{[s,t]} + D(s-)\delta_s - D(t-)\delta_t, \\ (D_{(s,t)})' &= (D')_{(s,t)} + D(s+)\delta_s - D(t-)\delta_t, \\ (D_{(s,t]})' &= (D')_{(s,t]} + D(s+)\delta_s - D(t+)\delta_t, \\ (D_{[s,t]})' &= (D')_{[s,t]} + D(s-)\delta_s - D(t+)\delta_t, \end{aligned}$$

where  $\delta_{\pm\infty} = 0$ .

The last part of this subsection considers matrices with piecewise-smoothly distributional entries and under which condition these matrices are invertible and how the inverse looks like.

**Definition 13 (Multiplication and invertibility of piecewise-smooth matrices)** For two matrices  $P \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times m}$ ,  $Q \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{m \times p}$ ,  $n, m, p \in \mathbb{N}$ , with piecewise-smoothly distributional entries the matrix product is defined in the standard way, i.e., for  $i = 1, \dots, n$  and  $j = 1, \dots, p$ ,

$$(PQ)_{ij} = \sum_{k=1}^m P_{ik}Q_{kj},$$

where  $M_{ij}$  denotes the  $(i, j)$ -entry of some matrix  $M$ . A square matrix  $M \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$ ,  $n \in \mathbb{N}$ , is called *invertible (over  $\mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ )* if, and only if, there exists a matrix  $M^{-1} \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  such that

$$MM^{-1} = M^{-1}M = I,$$

where  $I \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  is the (distributional) identity matrix given by

$$I_{ij} = \begin{cases} (\mathbb{1}_{\mathbb{R}})_{\mathbb{D}}, & i = j \\ 0, & i \neq j \end{cases}.$$

Note that no notational distinction between the matrices  $I \in \mathbb{R}^{n \times n}$ ,  $I \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$ , and  $I \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  is made.

**Proposition 14 (Invertibility of piecewise-smooth matrices)** Consider a piecewise-smoothly distributional matrix  $M \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$ ,  $n \in \mathbb{N}$ , with  $M_{\text{reg}}$  induced by  $M^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$ , i.e.  $M^{\text{reg}}_{\mathbb{D}} = M_{\text{reg}}$ . Then  $M$  is invertible if, and only if,  $M^{\text{reg}}$  is invertible over  $\mathcal{C}_{\text{pw}}^\infty$ , i.e. there exists  $P \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$  with  $M^{\text{reg}}(t)P(t) = P(t)M^{\text{reg}}(t) = I$  for all  $t \in \mathbb{R}$ .

If  $M$  is invertible, then the inverse is given by

$$M^{-1} = M_{\text{reg}}^{-1} - M_{\text{reg}}^{-1}M[\cdot]M_{\text{reg}}^{-1}, \quad \text{where } M_{\text{reg}}^{-1} := ((M^{\text{reg}})^{-1})_{\mathbb{D}}.$$

Note that for a matrix  $M^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$  the condition  $\det M^{\text{reg}}(t) \neq 0$  for all  $t \in \mathbb{R}$  is *not sufficient* for invertibility over  $\mathcal{C}_{\text{pw}}^\infty$ . Consider for example the  $1 \times 1$  matrix  $M^{\text{reg}}$  given by  $M^{\text{reg}}(t) = t$  on  $(-\infty, 0)$  and  $M^{\text{reg}}(t) = 1$  on  $[0, \infty)$  whose determinant is non-zero everywhere, but the inverse is  $(M^{\text{reg}})^{-1}(t) = 1/t$  on  $(-\infty, 0)$  and  $(M^{\text{reg}})^{-1}(t) = 1$  on  $[0, \infty)$  which is not a piecewise-smooth function because  $t \mapsto 1/t$  is not part of a globally smooth function as required by Definition 1.

### 3 Regularity of distributional DAEs

In this section, distributional DAEs of the form

$$E\dot{x} = Ax + f, \quad (11)$$

where  $E, A \in (\mathbb{D}_{\text{pwc}\infty})^{m \times n}$ ,  $m, n \in \mathbb{N}$ ,  $f \in (\mathbb{D}_{\text{pwc}\infty})^m$  and  $x \in (\mathbb{D}_{\text{pwc}\infty})^n$  are considered. Note that  $\dot{x} := x'$  is just used for traditional reasons.

#### 3.1 Definition of DAE-regularity

**Definition 15 (Initial trajectory problem (ITP))** Consider the distributional DAE (11), let  $x^0 \in (\mathbb{D}_{\text{pwc}\infty})^n$  and  $t_0 \in \mathbb{R}$ , then  $x \in (\mathbb{D}_{\text{pwc}\infty})^n$  is called a *solution of the initial trajectory problem (ITP)* (11) with initial trajectory  $x^0$  and initial time  $t_0$  if, and only if,

$$\begin{aligned} x_{(-\infty, t_0)} &= x_{(-\infty, t_0)}^0, \\ (E\dot{x})_{[t_0, \infty)} &= (Ax + f)_{[t_0, \infty)}. \end{aligned}$$

Now ITPs are used to define regularity of the matrix pair  $(E, A)$  in (11), in short,  $(E, A)$  is called regular if, and only if, every ITP is uniquely solvable. Note that the notion “regularity” is already used for distributions, therefore in the following the notion “DAE-regularity” is used to distinguish it from the distributional regularity. However, if the context is clear just “regularity” will be used.

**Definition 16 (DAE-regularity of  $(E, A)$ )** Consider the distributional DAE (11). The matrix pair  $(E, A)$  is called *DAE-regular* if, and only if, for all inhomogeneities  $f \in (\mathbb{D}_{\text{pwc}\infty})^m$ , for all initial trajectories  $x^0 \in (\mathbb{D}_{\text{pwc}\infty})^n$  and for all initial times  $t_0 \in \mathbb{R}$  the corresponding ITP has a unique solution.

Before formulating necessary and sufficient conditions for DAE-regularity, it is shown that regularity is invariant with respect to a certain system equivalence.

**Proposition 17 (Regularity and system equivalence)** Let  $S \in (\mathbb{D}_{\text{pwc}\infty})^{m \times m}$  and  $T \in (\mathbb{D}_{\text{pwc}\infty})^{n \times n}$  both be invertible over  $\mathbb{D}_{\text{pwc}\infty}$  and let  $(E, A)$  from (11) be DAE-regular. Then  $(\tilde{E}, \tilde{A}) := (SET, SAT - SET')$  is also DAE-regular.

In fact,  $\tilde{x}$  is a solution of the ITP (11) with  $(\tilde{E}, \tilde{A})$  and inhomogeneity  $\tilde{f} \in (\mathbb{D}_{\text{pwc}\infty})^m$ , initial trajectory  $\tilde{x}^0 \in (\mathbb{D}_{\text{pwc}\infty})^n$  and initial time  $t_0 \in \mathbb{R}$  if, and only if,  $x = T\tilde{x}$  is the solution of the ITP (11) with initial trajectory  $x^0 = T\tilde{x}^0$ , initial time  $t_0$  and inhomogeneity

$$f := S^{-1}\tilde{f}_{[t_0, \infty)} - S^{-1}[t_0] \left( \tilde{A}\tilde{x}^0 - \tilde{E}\dot{\tilde{x}}^0 \right)_{(-\infty, t_0)}.$$

### 3.2 Necessary conditions for DAE-regularity

**Theorem 18** ( $n = m$ ) *Consider the distributional DAE (11). If  $(E, A)$  is DAE-regular then  $n = m$ .*

This result is quite intuitive because, if  $n > m$  then there are more variables than equations, so the system is underdetermined, hence uniqueness of solutions can not be expected. If  $n < m$  then there are more equations than variables, hence the system is overdetermined and there exists inhomogeneities for which solutions do not exist. The next necessary conditions for regularity are more of technical nature.

**Theorem 19 (Derivative and impulse array)** *Consider the distributional DAE (11) with  $(E, A)$  DAE-regular.*

(i) *Define the derivative array of order  $p \in \mathbb{N}$  as a block matrix*

$$\mathcal{M}^p \in ((\mathbb{D}_{\text{pwC}^\infty})^{n \times n})^{p+1 \times p+2}$$

with, for  $i = 1, \dots, p+1$ ,  $j = 1, \dots, p+2$ ,

$$(\mathcal{M}^p)_{i,j} = \binom{i-1}{j-2} E^{(i-j+1)} - \binom{i-1}{j-1} A^{(i-j)},$$

with the convention that  $\binom{0}{0} = 1$  and  $\binom{n}{-k} = \binom{n}{n+k} = 0$  for  $k > 0$ ,  $n \in \mathbb{N}$ , i.e.

$$\mathcal{M}^p = \begin{bmatrix} -A & E & & & \\ -A' & E' - A & E & & \\ -A'' & E'' - 2A' & 2E' - A & E & \\ \vdots & \vdots & \vdots & \vdots & \ddots \\ -A^{(p)} & E^{(p)} - pA^{(p-1)} & pE^{(p-1)} - \binom{p}{2}A^{(p-2)} & \binom{p}{2}E^{(p-2)} - \binom{p}{3}A^{(p-3)} & \dots & E \end{bmatrix}$$

Then  $M^p(t+)$  and  $M^p(t-)$  have full row rank for all  $p \in \mathbb{N}$  and  $t \in \mathbb{N}$ .

(ii) *Define the impulse array of order  $(p, q)$ ,  $p, q \in \mathbb{N}$ , as a block matrix*

$$\mathcal{N}^{p,q} \in ((\mathbb{D}_{\text{pwC}^\infty})^{n \times n})^{p+1 \times q+1}$$

with, for  $i = 1, \dots, p+1$ ,  $j = 1, \dots, q+1$ ,

$$(\mathcal{N}^{p,q})_{i,j} = (-1)^{j-i} \left( \binom{j-1}{i-1} E^{(j-i)} + \binom{j-2}{i-1} A^{(j-i-1)} \right),$$

with the convention that  $\binom{-1}{k} = 0$  for all  $k \in \mathbb{N}$ , i.e.

$$\mathcal{N}^{p,q} = \begin{bmatrix} E - (E' + A) & E'' + A' \dots & (-1)^q (E^{(q)} + A^{(q-1)}) \\ -E & 2E' + A \dots & (-1)^{q-1} (qE^{(q-1)} + (q-1)A^{(q-2)}) \\ \dots & \dots & \vdots \\ (-1)^p E \dots & (-1)^{p-q} \left( \binom{q}{p} E^{q-p} + \binom{q-1}{p} A^{(q-p-1)} \right) \end{bmatrix}$$

Then for all  $p \in \mathbb{N}$  there exists  $q \in \mathbb{N}$  such that  $\mathcal{N}^{p,q}(t+)$  has full row rank for all  $t \in \mathbb{R}$ .

*Remark 20* Applying Theorem 19 to the time-invariant case both conditions reduces to the simple condition that all matrices

$$\begin{bmatrix} -A & E & & & & \\ & -A & E & & & \\ & & -A & E & & \\ & & & \ddots & \ddots & \\ & & & & -A & E \end{bmatrix}$$

have full row rank. Actually, this condition is equivalent to classical regularity of time-invariant DAEs [35].

### 3.3 Sufficient conditions for DAE-regularity

**Theorem 21 (Concatenation and additional impulses)** *Consider a family of distributional DAEs (11) with the corresponding matrix pairs  $(E_i, A_i)$ ,  $i \in \mathbb{Z}$ .*

(i) *If  $(E_0, A_0), (E_1, A_1)$  are DAE-regular, then*

$$(E, A) := (E_{0(-\infty, t_1)} + E_{1[t_1, \infty)}, A_{0(-\infty, t_1)} + A_{1[t_1, \infty)})$$

*is also DAE-regular for all  $t_1 \in \mathbb{R}$ .*

(ii) *If  $(E_i, A_i)$ ,  $i \in \mathbb{Z}$ , is DAE-regular and  $\{t_i \in \mathbb{R} \mid i \in \mathbb{Z}\}$  is a locally finite set, then*

$$(E, A) := \left( \sum_{i \in \mathbb{Z}} E_{i[t_i, t_{i+1})}, \sum_{i \in \mathbb{Z}} A_{i[t_i, t_{i+1})} \right)$$

*is also DAE-regular.*

(iii) *If  $(E_0, A_0)$  is DAE-regular, then*

$$(E, A) := (E_0 + E_1[t], A_0 + A_1[t])$$

*is also DAE-regular for all  $t \in \mathbb{R}$ .*

(iv) *If  $(E_0, A_0)$  is DAE-regular, then*

$$(E, A) := (E_0 + E_1[\cdot], A_0 + A_1[\cdot])$$

*is also DAE-regular.*

**Corollary 22** *Consider the distributional DAE (11), then  $(E, A)$  is DAE-regular if, and only if,  $(E_{\text{reg}}, A_{\text{reg}})$  is DAE-regular.*

*Remark 23* The Corollary 22 does *not* state that the impulses in  $E$  and  $A$  have no influence on the solutions, in fact, the proof of the Theorem 21 reveals that the impulsive parts of  $E$  and  $A$  are preserved in an altered inhomogeneity. In general, the presence of Dirac impulses and its derivatives in  $E$  and  $A$  yield solutions which might depend also on the derivatives of the initial trajectory.

Finally, some special distributional DAEs, namely distributional ordinary differential equations (ODEs) and pure distributional DAEs are studied and it turns out that these are DAE-regular.

**Theorem 24 (Regularity of ODEs and pure DAEs)** *Consider the distributional DAE (11). If  $(E, A) = (I, A)$  or  $(E, A) = (N, I)$ , where  $N \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  is such that  $N_{\text{reg}}$  is a strictly lower triangular matrix, then  $(E, A)$  is DAE-regular.*

*Remark 25 (Distributional ODEs)* The solution behavior of a distributional ODE  $\dot{x} = Ax + f$  differs significantly from the solution behavior of a classical ODE. Firstly, the solution of the ITP can depend on derivatives of the initial trajectory, so the “dimension” of the solution space can be larger than the size of the system. Secondly, it can be shown that for the free homogeneous distributional ODE  $\dot{x} = Ax$ , there exists, analogously as in the classical case, a fundamental solution  $\Phi_{t_0} \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$ , with  $t_0 \in \mathbb{R}$  such that  $A_{(-\infty, t_0)}[\cdot] = 0$ , i.e. every solution has the form  $x = \Phi_{t_0} x_0$  and  $x(t_0-) = x_0 \in \mathbb{R}^n$ . However, different to the classical case, the fundamental solution need not to be an invertible matrix. As an example, consider the distributional ODE  $\dot{x} = -\delta_0 x$  where all solutions are given by  $x = \mathbb{1}_{(-\infty, 0)} x_0$  for  $x_0 \in \mathbb{R}$ .

Finally the sufficient conditions can be summarized in the following way.

**Corollary 26 (Generalized Weierstraß form)** *Consider the distributional DAE (11). If there exist invertible matrices  $S, T \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$ , a locally finite set  $\{t_i \in \mathbb{R} \mid i \in \mathbb{Z}\}$ , a family of matrices  $J_i \in (\mathcal{C}_{\text{pw}}^\infty)^{n_i \times n_i}$ ,  $i \in \mathbb{Z}$ ,  $0 \leq n_i \leq n$  and a family of strictly lower triangular matrices  $N_i \in (\mathcal{C}_{\text{pw}}^\infty)^{(n-n_i) \times (n-n_i)}$ ,  $i \in \mathbb{Z}$  such that*

$$((SET)_{\text{reg}}, (SAT - SET')_{\text{reg}}) = \left( \sum_{i \in \mathbb{Z}} \begin{bmatrix} I & \\ & N_{i\mathbb{D}} \end{bmatrix}_{[t_i, t_{i+1})}, \sum_{i \in \mathbb{Z}} \begin{bmatrix} J_{i\mathbb{D}} & \\ & I \end{bmatrix}_{[t_i, t_{i+1})} \right),$$

then  $(E, A)$  is DAE-regular.

*Remark 27* For time-invariant DAEs, i.e.  $E, A \in \mathbb{R}^{n \times n}$ , the previous results show that DAE-regularity is identical to the classical regularity, defined by the condition  $\det(\lambda E - A) \in \mathbb{R}[\lambda] \setminus \{0\}$ .

## 4 Conclusion

To study time-varying DAEs of the form  $E\dot{x} = Ax + f$  with jumps in the coefficients, the space of piecewise-smooth distributions was introduced as a solution space. With this space it is also possible to allow for distributional entries in the coefficient matrices. The well known concept of regularity for classical DAEs (i.e. DAEs with constant coefficients) was generalized, necessary and sufficient conditions were given for the regularity of matrix pairs  $(E, A)$  with piecewise-smoothly distributional entries. It seems that even for



the classical time-varying case (i.e.  $E$  and  $A$  are smooth matrices) some of the conditions are new, in particular there is no general definition of regularity for time-varying DAEs. The presented framework is particularly suitable for studying switched DAEs, for example Theorem 21(ii) states that switching between regular systems yields a new regular system, and in general, without the presented framework it seems difficult to study switched DAEs at all.


Although regular DAEs play an important role, there are cases where non-regular DAEs also arise in applications, for example rectangular descriptions of systems. It seems that the “behavioral approach”, surveyed in [34], in combination with piecewise-smooth distributions as solutions will be a fruitful future research topic. Furthermore, for regular distributional DAEs, questions of control theory can be addressed and since the solution space as well as the control signal space are larger (they can include Dirac impulses) new methods and results are likely. Finally, the proposed distributional framework can be used to study reliability of linear networks, for example it is possible to study the situation that the failure of one component (which results in a new system description) induces impulsive solutions, which might destroy the system in reality.

## 5 Acknowledgements

I thank Achim Ilchmann for encouraging me to investigate this topic and also for all the valuable comments and suggestions for improvement. I would also like to thank Jan Willems for many helpful discussions. Furthermore, I would like to thank the anonymous reviewers for their constructive critics and for hinting to several interesting papers I wasn't aware of.

## Appendix 1: A simple electrical circuit example

Consider the simple circuit shown in Figure 1. In the circuit,  $C > 0$  is the capacity of the



**Fig. 1** A simple example circuit with a switch.

capacitor,  $R > 0$  is the resistance of the resistor and  $u : \mathbb{R} \rightarrow \mathbb{R}$  is the input voltage. The state variables are  $i_c$  and  $u_c$  which are the current through the capacitor and the voltage over the capacitor, respectively. Before time  $t = 0$ , the switch is on the right side, i.e. the capacitor is bypassed. At  $t = 0$ , the switch moves to the left and the capacitor starts charging. After

some time, the switch is moved back and the capacitor is bypassed again. The corresponding DAE reads as  $E \begin{pmatrix} u'_c \\ i'_c \end{pmatrix} = A \begin{pmatrix} u_c \\ i_c \end{pmatrix} + f$ , where

$$E = \begin{bmatrix} C & 0 \\ 0 & 0 \end{bmatrix}, \quad A(t) = \begin{cases} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{bmatrix} 0 & 1 \\ 1 & R \end{bmatrix}, & t \in [0, 1) \end{cases}, \quad f(t) = \begin{cases} \begin{pmatrix} 0 \\ 0 \end{pmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{pmatrix} 0 \\ -u(t) \end{pmatrix}, & t \in [0, 1). \end{cases}$$

Under the assumption that the input voltage satisfies  $u \equiv 1$ , the solution for the voltage  $u_c$  over the capacitor is given uniquely by

$$u_c(t) = \begin{cases} 0, & t \in \mathbb{R} \setminus [0, 1), \\ 1 - e^{-\frac{t}{RC}}, & t \in [0, 1). \end{cases}$$

Independently of the switch, the current  $i_c$  must fulfill the equation  $Cu'_c = i_c$ . *Since the solution of  $u_c$  has a jump, there is no classical solution for  $i_c$ .* However, if one allows for distributional solutions the equations are solvable and the (distributional) current  $i_c$  is given by

$$i_c = i_c^{\text{reg}}_{\mathbb{D}} + \underbrace{\left( e^{-\frac{1}{RC}} - 1 \right)}_{=u_c(1+) - u_c(1-)} \delta_1,$$

where  $i_c^{\text{reg}}$  is the regular part of the distribution given by

$$i_c^{\text{reg}}(t) = \begin{cases} 0, & t \in \mathbb{R} \setminus [0, 1), \\ \frac{1}{R} e^{-\frac{t}{RC}}, & t \in [0, 1) \end{cases}$$

and  $\delta_1$  is the Dirac impulse at  $t = 1$ .

Another way to find a solution is by transforming the DAE (locally) into the so called Weierstraß normal form [33] (see also [19, Thm. 2.12]) via

$$S(t) = \begin{cases} \begin{bmatrix} 0 & 1 \\ \frac{1}{C} & 0 \end{bmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{bmatrix} 1 & -\frac{1}{R} \\ 0 & \frac{1}{R} \end{bmatrix}, & t \in [0, 1), \end{cases} \quad T(t) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{bmatrix} \frac{1}{C} & 0 \\ -\frac{1}{RC} & 1 \end{bmatrix}, & t \in [0, 1). \end{cases}$$

The resulting DAE with  $(\tilde{E}, \tilde{A}) = (SET, SAT - SET')$  is then given by

$$\tilde{E}(t) = \begin{cases} \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, & t \in [0, 1) \end{cases}$$

and

$$\tilde{A}^{\text{reg}}(t) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{bmatrix} -\frac{1}{RC} & 0 \\ 0 & 1 \end{bmatrix}, & t \in [0, 1), \end{cases} \quad \tilde{A} = \tilde{A}^{\text{reg}}_{\mathbb{D}} + \begin{bmatrix} 0 & 0 \\ \frac{1-C}{C} & 0 \end{bmatrix} \delta_1.$$

Note that  $\tilde{A}$  now contains a Dirac impulse. Actually, the transformed DAE is not much simpler, but in general a DAE in Weierstraß form is easier to solve, here the transformation is just done to illustrate that impulses can occur in the coefficient matrices. The unique solution of  $\tilde{E}\dot{z} = \tilde{A}z + Sf$  is given by (again the input signal  $u$  is assumed to be constant and equal to one)

$$z^{\text{reg}}(t) = \begin{cases} \begin{pmatrix} 0 \\ 0 \end{pmatrix}, & t \in \mathbb{R} \setminus [0, 1), \\ \begin{pmatrix} C \left(1 - e^{-\frac{t}{RC}}\right) \\ \frac{1}{R} \end{pmatrix}, & t \in [0, 1), \end{cases} \quad z = z_{\mathbb{D}}^{\text{reg}} + \begin{pmatrix} 0 \\ e^{-\frac{1}{RC}} - 1 \end{pmatrix} \delta_1.$$

Now it is easy to verify that  $Tz$  is equal to the solution found above.

## Appendix 2: Proofs

### Proof of Proposition 2

By (3) it is, for every  $i \in \mathbb{Z}$ ,

$$\begin{aligned} (\mathbb{1}_{[t_i, t_{i+1})} f_i)_{\mathbb{D}}' &= (f_i (\mathbb{1}_{[t_i, t_{i+1})})_{\mathbb{D}})' \\ &= f_i' (\mathbb{1}_{[t_i, t_{i+1})})_{\mathbb{D}} + f_i (\mathbb{1}_{[t_i, \infty)} - \mathbb{1}_{[t_{i+1}, \infty)})_{\mathbb{D}}' \\ &= (\mathbb{1}_{[t_i, t_{i+1})} f_i')_{\mathbb{D}} + f_i \delta_{t_i} - f_i \delta_{t_{i+1}} \\ &= (\mathbb{1}_{[t_i, t_{i+1})} f_i')_{\mathbb{D}} + f_i(t_i) \delta_{t_i} - f_i(t_{i+1}) \delta_{t_{i+1}}. \end{aligned}$$

Now (7) follows from  $f_i(t_i) - f_{i-1}(t_i) = D(t_i+) - D(t_i-)$ . Finally, (2) implies that  $D[\cdot]'$  is again a locally finite sum of distributions with point support, hence  $D' \in \mathbb{D}_{\text{pw}C^\infty}$ .  $\square$

### Proof of Proposition 3

As already mentioned, every distribution  $D \in \mathbb{D}$  has a distributional antiderivative and all antiderivatives only differ by a constant. It is first shown, that every distributional antiderivative  $H$  of a piecewise-smooth distribution  $D \in \mathbb{D}_{\text{pw}C^\infty}$  is a piecewise-smooth distribution. Consider the representation  $D = f_{\mathbb{D}} + \sum_{t \in T} D_t \in \mathbb{D}_{\text{pw}C^\infty}$  as in Definition 1. Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be an antiderivative of  $f$ , then  $g \in C_{\text{pw}}^\infty$ . For a fixed  $t \in T$  and by (2),  $D_t$  can be written as

$$D_t = \sum_{i=0}^{n_t} a_t^i \delta_t^{(i)},$$

where  $n_t \in \mathbb{N}$  and  $a_t^0, \dots, a_t^{n_t} \in \mathbb{R}$ . Clearly, one antiderivative of  $D_t$  is given by

$$a_t^0 (\mathbb{1}_{[t, \infty)})_{\mathbb{D}} + \sum_{i=1}^{n_t} a_t^i \delta_t^{(i-1)}.$$

Now let

$$h = g + \sum_{t \in T} a_t^0 \mathbb{1}_{[t, \infty)} \in C_{\text{pw}}^\infty$$

and, for  $t \in T$ ,

$$\tilde{D}_t = \sum_{i=1}^{n_t} a_t^i \delta_t^{(i-1)},$$

then  $H_1 = h_{\mathbb{D}} + \sum_{t \in T} \tilde{D}_t \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$  is a distributional antiderivative of  $D$ . Since all other antiderivatives only differ by a constant, all antiderivatives of  $D$  are piecewise-smooth distributions. Let

$$H = H_1 - H_1(t_0-) \mathbb{1}_{\mathbb{R}\mathbb{D}}$$

then  $H$  is the only distributional antiderivative with the property  $H(t_0-) = 0$ . □

### Proof of Theorem 4

First observe that in (9) locally finiteness of  $T_F$  and  $T_G$  together with (5) ensures that indeed  $\sum_{t \in T_F} F[t]g_{\mathbb{D}} = F[\cdot]g_{\mathbb{D}}$  and  $\sum_{t \in T_G} f_{\mathbb{D}}G[t] = f_{\mathbb{D}}G[\cdot]$ .

- (i) Clearly,  $fg \in \mathcal{C}_{\text{pw}}^\infty$ , hence it remains to show that  $\sum_{t \in T_F} F[t]g_{\mathbb{D}}$  and  $\sum_{t \in T_G} f_{\mathbb{D}}G[t]$  are piecewise-smooth distributions. From the definition it follows that

$$\begin{aligned} \forall t \in T_F : \quad \text{supp}(F[t]g_{\mathbb{D}}) &\subseteq \{t\} \\ \forall t \in T_G : \quad \text{supp}(f_{\mathbb{D}}G[t]) &\subseteq \{t\} \end{aligned}$$

which shows that  $FG \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ .

- (ii) This directly follows from the definition since  $f_{\mathbb{D}}[\cdot] = 0 = g_{\mathbb{D}}[\cdot]$ .  
 (iii) Simple calculations show the validity of these properties.  
 (iv) This part of the proof consists of four steps.

*Step 1:* It is shown that  $(fg)_{\mathbb{D}}' = f_{\mathbb{D}}'g_{\mathbb{D}} + f_{\mathbb{D}}g_{\mathbb{D}}'$ .

Let  $f = \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i$  and  $g = \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} g_i$  with  $f_i, g_i \in \mathcal{C}^\infty$ ,  $i \in \mathbb{Z}$ , and locally finite  $\{t_i \in \mathbb{R} \mid i \in \mathbb{Z}\}$ . Note that both representations use the same intervals, but this is no restriction of generality. Then, by (7),

$$(fg)_{\mathbb{D}}' = \left( \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} (f_i g_i)' \right)_{\mathbb{D}} + \sum_{i \in \mathbb{Z}} (f(t_i+)g(t_i+) - f(t_i-)g(t_i-)) \delta_{t_i},$$

and

$$\begin{aligned} f_{\mathbb{D}}'g_{\mathbb{D}} &= \left( \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i' g_i \right)_{\mathbb{D}} + \sum_{i \in \mathbb{Z}} (f(t_i+) - f(t_i-)) \delta_{t_i} g_{\mathbb{D}}, \\ f_{\mathbb{D}}g_{\mathbb{D}}' &= \left( \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i g_i' \right)_{\mathbb{D}} + \sum_{i \in \mathbb{Z}} f_{\mathbb{D}}(g(t_i+) - g(t_i-)) \delta_{t_i}. \end{aligned}$$

Since  $(f_i g_i)' = f_i' g_i + f_i g_i'$ ,  $\delta_{t_i} g_{\mathbb{D}} = g(t_i-) \delta_{t_i}$ , and  $f_{\mathbb{D}} \delta_{t_i} = f(t_i+) \delta_{t_i}$  for all  $i \in \mathbb{Z}$ , the assertion of Step 1 is shown.

*Step 2:* It is shown that  $(F[t]g_{\mathbb{D}})' = F[t]'g_{\mathbb{D}} + F[t]g_{\mathbb{D}}'$  for all  $t \in T_F$ .

Since  $F[t]$  is by (2) a finite sum of a Dirac impulse and its derivatives, it suffices to consider the case  $F[t] = \delta_t^{(n)}$  for some  $n \in \mathbb{N}$ . Now the assertion follows directly from (8).

*Step 3:* It is shown that  $(f_{\mathbb{D}}G[t])' = f_{\mathbb{D}}'G[t] + f_{\mathbb{D}}G[t]'$  for all  $t \in T_G$ .

As in Step 2 it suffices to consider  $G[t] = \delta_t^{(n)}$  for some  $n \in \mathbb{N}$ . Now the assertion follows again from (8).

*Step 4:*  $(FG)' = F'G + FG'$  is shown.

Since  $T_F$  and  $T_G$  are locally finite it follows from Step 2 and 3 that

$$\begin{aligned} (F[\cdot]g_{\mathbb{D}})' &= F[\cdot]'g_{\mathbb{D}} + F[\cdot]g_{\mathbb{D}}' \\ (f_{\mathbb{D}}G[\cdot])' &= f_{\mathbb{D}}'G[\cdot] + f_{\mathbb{D}}G[\cdot]' \end{aligned}$$

Expanding the products yields

$$(FG)' = (fg)_{\mathbb{D}}' + (F[\cdot]g_{\mathbb{D}})' + (f_{\mathbb{D}}G[\cdot])'$$

and

$$\begin{aligned} F'G + FG' &= (f_{\mathbb{D}})'g_{\mathbb{D}} + (f_{\mathbb{D}})'G[\cdot] + F[\cdot]'g_{\mathbb{D}} + F[\cdot]'G[\cdot] \\ &\quad + f_{\mathbb{D}}g_{\mathbb{D}}' + f_{\mathbb{D}}G[\cdot]' + F[\cdot]g_{\mathbb{D}}' + F[\cdot]G[\cdot]'. \end{aligned}$$

Hence

$$(FG)' - (F'G + FG') = (fg)_{\mathbb{D}}' - (f_{\mathbb{D}})'g_{\mathbb{D}} - (f_{\mathbb{D}})'G[\cdot] - F[\cdot]'g_{\mathbb{D}} - F[\cdot]'G[\cdot].$$

The regular parts of  $F[\cdot]'$  and  $G[\cdot]'$  are zero, hence  $F[\cdot]'G[\cdot] = 0$  and  $F[\cdot]G[\cdot]' = 0$ . Using the equality from Step 1 now yields the assertion.

(v) This follows from

$$\begin{aligned} \text{supp } FG &= \text{supp } (f_{\mathbb{D}}g_{\mathbb{D}} + f_{\mathbb{D}}G[\cdot] + F[\cdot]g_{\mathbb{D}}) \\ &\subseteq (\text{supp } f_{\mathbb{D}} \cap \text{supp } g_{\mathbb{D}}) \cup (\text{supp } f_{\mathbb{D}} \cap \text{supp } G[\cdot]) \\ &\quad \cup (\text{supp } F[\cdot] \cap \text{supp } g_{\mathbb{D}}) \cup (\text{supp } F[\cdot] \cap \text{supp } G[\cdot]) \\ &= (\text{supp } f_{\mathbb{D}} \cup \text{supp } F[\cdot]) \cap (\text{supp } g_{\mathbb{D}} \cup \text{supp } G[\cdot]) \\ &= \text{supp } F \cap \text{supp } G. \end{aligned}$$

□

## Proof of Proposition 11

Let  $M \subseteq \mathbb{R}$  be one of the four intervals with boundaries  $s$  and  $t$ , then by linearity of the restriction

$$(FG)_M = (F_{\text{reg}}G_{\text{reg}})_M + (F_{\text{reg}}G[\cdot])_M + (F[\cdot]G_{\text{reg}})_M.$$

First observe that  $(F_{\text{reg}}G_{\text{reg}})_M = (F_{\text{reg}})_M(G_{\text{reg}})_M$ . Furthermore,

$$(F_{\text{reg}}G[\cdot])_M = ((F_{\text{reg}})_M G[\cdot])_M + ((F_{\text{reg}})_{\mathbb{R} \setminus M} G[\cdot])_M + (F_{\text{reg}}G[\cdot]_{\mathbb{R} \setminus M})_M,$$

where the term  $(F_{\text{reg}}G[\cdot]_{\mathbb{R} \setminus M})_M$  is zero, because  $F_{\text{reg}}G[\cdot]_{\mathbb{R} \setminus M}$  is a distribution with zero regular part and whose support is a locally finite set contained in  $\mathbb{R} \setminus M$ , hence the restriction to  $M$  is zero by definition. Since the support of  $(F_{\text{reg}})_M G[\cdot]_M$  is a locally finite set and is contained within  $M$  the outer restriction does not change it. Finally, the support of  $(F_{\text{reg}})_{\mathbb{R} \setminus M} G[\cdot]_M$  is also a locally finite set and is contained in  $\{s, t\}$ , hence, if  $s < t$ ,

$$(F_{\text{reg}}G[\cdot])_M = (F_{\text{reg}})_M G[\cdot]_M + (F_{\text{reg}})_{\mathbb{R} \setminus M} (G[s] + G[t])_M$$

Analogously,

$$(F[\cdot]G_{\text{reg}})_M = F[\cdot]_M (G_{\text{reg}})_M + (F[s] + F[t])_M (G_{\text{reg}})_{\mathbb{R} \setminus M}.$$

Now let  $M = (s, t)$ , then  $(G[s] + G[t])_M = 0 = (F[s] + F[t])_M$ , hence the assertion is shown in this case. For  $M = [s, t)$  it is  $(G[s] + G[t])_M = G[s]$  and  $(F[s] + F[t])_M = F[s]$ . From (10) it follows that the term  $(F_{\text{reg}})_{\mathbb{R} \setminus M} G[s]$  depends only on the value  $((F_{\text{reg}})_{\mathbb{R} \setminus M})^{(i)}(s+)$ ,  $i \in \mathbb{N}$ , which is zero for all  $i \in \mathbb{N}$ , hence  $(F_{\text{reg}})_{\mathbb{R} \setminus M} G[s] = 0$ . Also from (10) it follows that  $F[s](G_{\text{reg}})_{\mathbb{R} \setminus M} = F[s]G_{(s-\varepsilon, s)}$  for any  $\varepsilon > 0$ . This shows the assertion for  $M = [s, t)$ . Analogous arguments show the validity of the assertions for  $M = (s, t]$  and  $M = [s, t]$ .

If  $s = t$ , then

$$\begin{aligned} (FG)_{[s, t]} &= (FG)[s] = (F_{\text{reg}}G[\cdot])[s] + (F[\cdot]G_{\text{reg}})[s] \\ &= F_{\text{reg}}G[s] + F[s]G_{\text{reg}} \\ &= \underbrace{F[s]G[s]}_{=0} + F_{\text{reg}(s, s+\varepsilon)}G[s] + F[s]G_{\text{reg}(s-\varepsilon, s)} \\ &= F_{[s, t]}G_{[s, t]} + F_{(t, t+\varepsilon)}G[s] + F[s]G_{(s-\varepsilon, s)} \end{aligned}$$

□

### Proof of Proposition 12

Let  $D_{\text{reg}} = \left( \sum_{i \in \mathbb{Z}} \mathbb{1}_{[t_i, t_{i+1})} f_i \right)_{\mathbb{D}}$  for some locally finite set  $\{ t_i \in \mathbb{R} \mid i \in \mathbb{Z} \}$  and  $f_i \in \mathcal{C}^\infty$ ,  $i \in \mathbb{Z}$ . Assume, without restriction, that  $s, t \in \{ t_i \mid i \in \mathbb{Z} \}$ . From (7) follows

$$\begin{aligned} (D_{[s,t]})' - (D')_{[s,t]} &= \sum_{i \in \mathbb{Z}} (D_{[s,t]}(t_i+) - D_{[s,t]}(t_i-)) \delta_{t_i} \\ &\quad - \left( \sum_{i \in \mathbb{Z}} (D(t_i+) - D(t_i-)) \delta_{t_i} \right)_{[s,t]} \\ &= (D_{[s,t]}(s+) - D_{[s,t]}(s-)) \delta_s + (D_{[s,t]}(t+) - D_{[s,t]}(t-)) \delta_t \\ &\quad - (D(s+) - D(s-)) \delta_s \\ &= -D(t-)\delta_t + D(s-)\delta_s. \end{aligned}$$

This shows the first formula. Since  $D[\tau]' = D'[\tau] - (D(\tau+) - D(\tau-))\delta_\tau$  for all  $\tau \in \mathbb{R}$  the other three formulae follow easily.  $\square$

### Proof of Proposition 14

If  $M^{\text{reg}}$  is invertible over  $\mathcal{C}_{\text{pw}}^\infty$  then

$$\begin{aligned} MM^{-1} &= (M_{\text{reg}} + M[\cdot]) (M_{\text{reg}}^{-1} - M_{\text{reg}}^{-1}M[\cdot]M_{\text{reg}}^{-1}) \\ &= \underbrace{M_{\text{reg}}M_{\text{reg}}^{-1}}_{=I} - \underbrace{M_{\text{reg}}M_{\text{reg}}^{-1}M[\cdot]M_{\text{reg}}^{-1} + M[\cdot]M_{\text{reg}}^{-1}}_{=0} - \underbrace{M[\cdot]M_{\text{reg}}^{-1}M[\cdot]M_{\text{reg}}^{-1}}_{=0}, \end{aligned}$$

where the last zero follows from the fact the product of two piecewise-smooth distributions with zero regular part is zero. An analogous calculation shows  $M^{-1}M = I$ . Hence sufficiency is shown.

Now assume that  $M$  is invertible over  $\mathbb{D}_{\text{pw}\mathcal{C}^\infty}$ , i.e. there exists a matrix  $M^{-1} \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  such that  $MM^{-1} = I$ . Let  $M^{-1} = (M^{-1})_{\text{reg}} + M^{-1}[\cdot]$ , then

$$\begin{aligned} I &= MM^{-1} = (M_{\text{reg}} + M[\cdot])((M^{-1})_{\text{reg}} + M^{-1}[\cdot]) \\ &= M_{\text{reg}}(M^{-1})_{\text{reg}} + \underbrace{M_{\text{reg}}M^{-1}[\cdot] + M[\cdot](M^{-1})_{\text{reg}}}_{=:H}. \end{aligned}$$

Since  $H[\cdot] = H$  and  $I[\cdot] = 0$ , it follows that  $H$  must be zero. This implies

$$I = M_{\text{reg}}(M^{-1})_{\text{reg}} = (M^{\text{reg}}(M^{-1})^{\text{reg}})_{\mathbb{D}}$$

where  $(M^{-1})^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$  is such that  $(M^{-1})_{\text{reg}} = (M^{-1})^{\text{reg}}_{\mathbb{D}}$ . Hence  $M^{\text{reg}}$  is invertible over  $\mathcal{C}_{\text{pw}}^\infty$  with inverse  $(M^{-1})^{\text{reg}}$ . Finally, from  $H = 0$  and the invertibility of  $M_{\text{reg}}$  it follows that

$$M^{-1}[\cdot] = -(M_{\text{reg}})^{-1}M[\cdot](M^{-1})_{\text{reg}} = -M_{\text{reg}}^{-1}M[\cdot]M_{\text{reg}}^{-1},$$

hence  $M^{-1}$  is unique.  $\square$

### Proof of Proposition 17

It will be shown that every ITP

$$\tilde{E}\tilde{x} = \tilde{A}\tilde{x} + \tilde{f}, \quad \tilde{x}_{(-\infty, t_0)} = \tilde{x}_{(-\infty, t_0)}^0,$$

with  $t_0 \in \mathbb{R}$ ,  $\tilde{x}^0 \in \mathbb{D}_{\text{pwc}\infty}^n$ ,  $\tilde{f} \in \mathbb{D}_{\text{pwc}\infty}^m$ , has a unique solution.

*Step 1:* Existence of a solution.

Let  $x$  be the solution of the ITP

$$E\dot{x} = Ax + f, \quad x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0,$$

where

$$f = S^{-1}\tilde{f}_{[t_0, \infty)} - S^{-1}[t_0] \left( \tilde{A}\tilde{x}^0 - \tilde{E}\dot{\tilde{x}}^0 \right)_{(-\infty, t_0)}$$

and  $x^0 = T\tilde{x}^0$ . It will be shown that  $\tilde{x} := T^{-1}x$  is the desired solution. First observe that, by Proposition 11,

$$\tilde{x}_{(-\infty, t_0)} = (T^{-1}x)_{(-\infty, t_0)} = T_{(-\infty, t_0)}^{-1}x_{(-\infty, t_0)}^0 = T_{(-\infty, t_0)}^{-1}(T\tilde{x}^0)_{(-\infty, t_0)} = \tilde{x}_{(-\infty, 0)}^0.$$

Hence it remains to show that

$$(\tilde{E}\dot{\tilde{x}})_{[t_0, \infty)} = (\tilde{A}\tilde{x})_{[t_0, \infty)} + \tilde{f}_{[t_0, \infty)},$$

which is equivalent to

$$S^{-1}(\tilde{E}\dot{\tilde{x}})_{[t_0, \infty)} = S^{-1}(\tilde{A}\tilde{x})_{[t_0, \infty)} + S^{-1}\tilde{f}_{[t_0, \infty)}.$$

Note that from Proposition 11 and Property (R4) it follows that, for any  $M \in (\mathbb{D}_{\text{pwc}\infty})^{m \times h}$  and  $h = 1$  or  $h = n$ ,

$$\begin{aligned} S^{-1}M_{[t_0, \infty)} &= (S^{-1}M_{[t_0, \infty)})_{(-\infty, t_0)} + (S^{-1}M_{[t_0, \infty)})_{[t_0, \infty)} \\ &= 0 + S_{[t_0, \infty)}^{-1}M_{[t_0, \infty)} \\ &= (S^{-1}M)_{[t_0, \infty)} - S^{-1}[t_0]M_{(-\infty, t_0)}. \end{aligned}$$

Hence  $\tilde{x}$  must fulfill

$$\begin{aligned} &(S^{-1}\tilde{E}\dot{\tilde{x}})_{[t_0, \infty)} - S^{-1}[t_0](\tilde{E}\dot{\tilde{x}})_{(-\infty, t_0)} \\ &= (S^{-1}\tilde{A}\tilde{x})_{[t_0, \infty)} - S^{-1}[t_0](\tilde{A}\tilde{x})_{(-\infty, t_0)} + (S^{-1}\tilde{f})_{[t_0, \infty)} - S^{-1}[t_0]\tilde{f}_{(-\infty, t_0)}. \end{aligned}$$

From  $0 = (T^{-1}T)' = (T^{-1})'T + T^{-1}T'$  it follows, that

$$(T^{-1})' = -T^{-1}T'T^{-1},$$

hence

$$S^{-1}\tilde{E}\dot{\tilde{x}} = S^{-1}SET(T^{-1}x)' = E\dot{x} - ET'T^{-1}x$$

and

$$S^{-1}\tilde{A}\tilde{x} = S^{-1}(SAT - SET')T^{-1}x = Ax - ET'T^{-1}x.$$

Since, by assumption,  $(E\dot{x})_{[t_0, \infty)} = (Ax)_{[t_0, \infty)} + f_{[t_0, \infty)}$ , it remains to show that

$$f_{[t_0, \infty)} = S^{-1}[t_0](\tilde{E}\dot{\tilde{x}})_{(-\infty, t_0)} - S^{-1}[t_0](\tilde{A}\tilde{x})_{(-\infty, t_0)} + (S^{-1}\tilde{f})_{[t_0, \infty)} - S^{-1}[t_0]\tilde{f}_{(-\infty, t_0)}.$$

Together with Proposition 11 and Proposition 12 this follows from

$$\begin{aligned} (\tilde{E}\dot{\tilde{x}})_{(-\infty, t_0)} &= \tilde{E}_{(\infty, t_0)}\dot{\tilde{x}}_{(\infty, t_0)} \\ &= \tilde{E}_{(\infty, t_0)} \left( (\tilde{x}_{(-\infty, t_0)})' + \tilde{x}(t_0-)\delta_{t_0} \right) \\ &= \tilde{E}_{(\infty, t_0)} \left( (\tilde{x}_{(-\infty, t_0)}^0)' + \tilde{x}^0(t_0-)\delta_{t_0} \right) \\ &= \tilde{E}_{(\infty, t_0)}\dot{\tilde{x}}_{(\infty, t_0)}^0 = (\tilde{E}\dot{\tilde{x}}^0)_{(-\infty, t_0)}, \\ (\tilde{A}\tilde{x})_{(-\infty, t_0)} &= (\tilde{A}\tilde{x}^0)_{(-\infty, t_0)}, \\ (S^{-1}\tilde{f})_{[t_0, \infty)} - S^{-1}[t_0]\tilde{f}_{(-\infty, t_0)} &= S_{[t_0, \infty)}^{-1}\tilde{f}_{[t_0, \infty)} = S^{-1}\tilde{f}_{[t_0, \infty)}, \end{aligned}$$

and the definition of  $f$ .

*Step 2:* Uniqueness of a solution.

Let  $\tilde{x}_1$  and  $\tilde{x}_2$  be two solutions of the ITP

$$\tilde{E}\dot{\tilde{x}} = \tilde{A}\tilde{x} + \tilde{f}, \quad \tilde{x}_{(-\infty, t_0)} = \tilde{x}_{(-\infty, t_0)}^0$$

for some  $t_0 \in \mathbb{R}$ ,  $\tilde{x}^0 \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}^n$ ,  $\tilde{f} \in \mathbb{D}_{\text{pw}\mathcal{C}^\infty}^m$ . Then  $\tilde{z} := \tilde{x}_1 - \tilde{x}_2$  is a solution of the ITP

$$\tilde{E}\dot{\tilde{z}} = \tilde{A}\tilde{z}, \quad \tilde{z}_{(-\infty, t_0)} = 0.$$

It will be shown that  $z = T\tilde{z}$  is a solution of the ITP

$$E\dot{z} = Az, \quad z_{(-\infty, t_0)} = 0,$$

it then follows from the DAE-regularity of  $(E, A)$  that  $z = 0$ , hence  $\tilde{z} = 0$  and the uniqueness of solutions is shown.

Clearly,  $z_{(-\infty, t_0)} = 0$ , hence it remains to show that  $(E\dot{z})_{[t_0, \infty)} = (Az)_{[t_0, \infty)}$ . It is, by Proposition 11,

$$\begin{aligned} 0 &= (\tilde{E}\dot{\tilde{z}})_{[t_0, \infty)} - (\tilde{A}\tilde{z})_{[t_0, \infty)} \\ &= S_{[t_0, \infty)}(E\dot{z} - Az)_{[t_0, \infty)} + S[t_0](E\dot{z} - Az)_{(-\infty, t_0)} \\ &= S(E\dot{z} - Az)_{[t_0, \infty)} + 0, \end{aligned}$$

hence  $(E\dot{z})_{[t_0, \infty)} = (Az)_{[t_0, \infty)}$ . □

## Proof of Theorem 18

*Step 1:*  $m \leq n$

Seeking a contradiction assume  $m > n$ . Let  $E = E^{\text{reg}}_{\mathbb{D}} + E[\cdot]$ , where  $E^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{m \times n}$ . Let  $r_E : \mathbb{R} \rightarrow \mathbb{N}$ ,  $t \mapsto r_E(t) := \text{rk } E^{\text{reg}}(t)$ , then there exists an open Interval  $J \subseteq \mathbb{R}$  such that  $r_E$  is constant on  $J$  (see e.g. [19, Thm. 3.25]) and  $E^{\text{reg}}|_J$  is smooth. Let  $r := r(t)$  for some  $t \in J$ , then  $r \leq n < m$ . In particular, there exists an invertible  $S \in (\mathcal{C}^\infty(J \rightarrow \mathbb{R}))^{m \times m}$  such that

$$SE^{\text{reg}}|_J = \begin{bmatrix} \tilde{E} \\ 0_{(m-n-r) \times n} \\ 0_{(m-n) \times n} \end{bmatrix} =: \begin{bmatrix} \hat{E} \\ 0_{(m-n) \times n} \end{bmatrix}$$

for some  $\tilde{E} \in (\mathcal{C}^\infty(J \rightarrow \mathbb{R}))^{r \times n}$  (Doležal's Theorem, [4]) and corresponding  $\hat{E} \in (\mathcal{C}^\infty(J \rightarrow \mathbb{R}))^{n \times n}$ . Without restriction, it can be assumed that  $\inf_{t \in J} \det S(t) > 0$  and  $E_J[\cdot] = A_J[\cdot] = 0$  (if these conditions are not fulfilled a reduction of the size of the open interval  $J$  yields these properties). Hence it is possible to extend the matrix function  $S$  to the whole time interval  $\mathbb{R}$  such that  $S \in (\mathcal{C}^\infty)^{m \times m}$  and  $S^{-1} \in (\mathcal{C}^\infty)^{m \times m}$  exists and

$$SE = \begin{bmatrix} E_1 \\ E_2 \end{bmatrix}, \quad SA = \begin{bmatrix} A_1 \\ A_2 \end{bmatrix},$$

where  $E_1, A_1 \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{n \times n}$  and  $(E_2)_J = 0$ . By Proposition 17 the pair  $(SE, SA)$  is still DAE-regular, in particular the DAE must have a local solution on the interval  $J$  for all inhomogeneities. Let  $A_2 = A_2^{\text{reg}}_{\mathbb{D}} + A_2[\cdot]$  for some  $A_2^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{(m-n) \times (m-n)}$ . It now follows that  $A_2^{\text{reg}}|_J$  must have full row rank, because otherwise there would exist  $t \in J$  and an invertible matrix  $M \in \mathbb{R}^{m \times m}$  such that the last row of  $MSA(t+)$  and of  $MSE(t+)$  is zero, hence for every inhomogeneity  $f$  with  $f(t+) \neq 0$  any ITP with  $t_0 \leq t$  would not have a solution. Firstly, this implies  $m \leq 2n$ . Secondly, by Doležal's Theorem there exists a matrix function  $T \in \mathcal{C}^\infty(J \rightarrow \mathbb{R})^{n \times n}$  such that

$$A_2T = [0 \ I] \text{ on } J$$



and it is possible to extend  $T$  on the whole axis, such that  $T \in (\mathcal{C}^\infty)^{n \times n}$  with  $T^{-1} \in (\mathcal{C}^\infty)^{n \times n}$  (possible by reducing the size of  $J$ ). Now let

$$(\tilde{E}, \tilde{A}) := (SET, SAT - SET') = \left( \begin{bmatrix} E_{11} & E_{12} \\ E_{21} & E_{22} \end{bmatrix}, \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \right),$$

then  $(\tilde{E}, \tilde{A})$  is DAE-regular by Proposition 17 and

$$E_{21J} = 0, E_{22J} = 0, A_{21J} = 0, A_{22J} = I_J,$$

furthermore, the size of  $E_{11}$  is  $n \times (2n - m)$ . Since  $m > n$  was assumed it follows that  $E_{11}$  has a strictly smaller size than  $E$  and has more rows than columns. On the interval  $J$  the system  $(\tilde{E}, \tilde{A})$  reads as

$$\begin{aligned} E_{11}\dot{z}_1 + E_{12}\dot{z}_2 &= A_{11}z_1 + A_{12}z_2 + f_1, \\ 0 &= z_2 + f_2, \end{aligned}$$

where  $f_1 \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^n$ ,  $f_2 \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{m-n}$ ,  $z_1 \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{2n-m}$ ,  $z_2 \in (\mathbb{D}_{\text{pw}\mathcal{C}^\infty})^{m-n}$ . If  $2m = n$ , then the above equation is equivalent to

$$f_1 = A_{12}f_2 - E_{12}f_2' \quad \text{on } J,$$

hence it is only solvable on the interval  $J$  if the inhomogeneity is not chosen arbitrarily. Hence  $2m > n$ . In this case, substituting  $z_2$  in the above equation by  $f_2$  yields that the system  $(E, A)$  has a local solution on the interval  $J$  if, and only if, the system  $(E_{11}, A_{11})$  has a local solution on the interval  $J$ . Let  $(E^0, A^0) := (E, A)$  and  $(E^1, A^1) := (E_{11}, A_{11})$  with size  $m^0 \times n^0 := m \times n$  and  $m^1 \times n^1 := n \times (2n - m)$ . It is now possible to repeat the above arguments to get a sequence of matrix-pairs  $(E^i, A^i)$ ,  $i \in \mathbb{N}$ , with strictly decreasing size  $m^i \times n^i$  such that  $0 \leq m^i < m^{i-1}$  and  $0 \leq n^i < n^{i-1}$ . Clearly, this is a contradiction. *Step 2:  $n \leq m$*

Seeking a contradiction assume  $n > m$ . With analogous arguments as in the first step it is possible to find an open interval  $J \subseteq \mathbb{R}$  and invertible matrices  $S \in (\mathcal{C}^\infty)^{n \times n}$ ,  $T \in (\mathcal{C}^\infty)^{m \times m}$  such that

$$(\tilde{E}, \tilde{A}) := (SET, SAT - SET') = \left( \begin{bmatrix} E_{11} & E_{12} \\ E_{21} & E_{22} \end{bmatrix}, \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \right)$$

is DAE-regular and

$$E_{12J} = 0, E_{22J} = 0, A_{12J} = 0, A_{22J} = I_J.$$

To get this result it was used that  $S^{-1} \begin{bmatrix} A_{12} \\ A_{22} \end{bmatrix}$  must have full column rank on  $J$ , because otherwise there would exist a component of the solution vector  $x$  which does not “appear” in the DAE on  $J$  and hence the ITP could not have a unique solution. The size of  $E_{11}$  is  $(2m - n) \times n$ , from which follows that  $2m \geq n$ . The system  $(\tilde{E}, \tilde{A})$  restricted to  $J$  reads as

$$\begin{aligned} E_{11}\dot{z}_1 &= A_{11}z_1 + f_1 \\ E_{21}\dot{z}_1 &= A_{21}z_1 + z_2 + f_2. \end{aligned}$$

If  $2m = n$  then only the equation

$$z_2 = A_{21}z_1 - E_{21}\dot{z}_1 + f_2$$

remains, which is clearly not uniquely solvable (with a given initial trajectory) on the interval  $J$ , because  $z_1$  can be altered on  $J$  and together with the corresponding  $z_2$  it is still a solution of the same ITP. Hence  $2m > n$ . Similar as in the first step it is now again possible to construct a sequence of systems  $(E^i, A^i)$ ,  $i \in \mathbb{N}$ , with strictly decreasing size  $m^i \times n^i$  such that  $m^i > n^i$  and  $0 < m^i < m^{i-1}$ . This is a contradiction.  $\square$

## Proof of Theorem 19

(i) Taking successively the derivative of the equation  $E\dot{x} = Ax + f$  yields

$$\begin{aligned} E\dot{x} - Ax &= f \\ E\ddot{x} + (E' - A)\dot{x} - A'x &= f' \\ E\ddot{\ddot{x}} + (2E' - A)\ddot{x} + (E'' - 2A')\dot{x} - A''x &= f'' \\ &\vdots \end{aligned}$$

and it follows inductively that, for all  $p \in \mathbb{N}$ ,

$$\mathcal{M}^p \begin{pmatrix} x \\ \dot{x} \\ \vdots \\ x^{(p)} \\ x^{(p+1)} \end{pmatrix} = \begin{pmatrix} f \\ f' \\ \vdots \\ f^{(p)} \end{pmatrix}$$

and, in particular, for all  $t \in \mathbb{R}$ ,

$$\mathcal{M}^p(t_{\pm}) \begin{pmatrix} x(t_{\pm}) \\ \dot{x}(t_{\pm}) \\ \vdots \\ x^{(p)}(t_{\pm}) \\ x^{(p+1)}(t_{\pm}) \end{pmatrix} = \begin{pmatrix} f(t_{\pm}) \\ f'(t_{\pm}) \\ \vdots \\ f^{(p)}(t_{\pm}) \end{pmatrix}$$

Since  $(E, A)$  is assumed to be DAE-regular there exists a solution for any given right-hand side, hence  $\mathcal{M}^p(t_+)$  and  $\mathcal{M}^p(t_-)$  must both have full row rank.

(ii) For a fixed  $t_0 \in \mathbb{R}$  consider the impulsive part of the DAE (11) at  $t_0$ :

$$(E\dot{x})[t_0] = (Ax + f)[t_0]$$

or, equivalently,

$$E_{(t_0, \infty)}\dot{x}[t_0] - A_{(t_0, \infty)}x[t_0] = A[t_0]x_{(-\infty, t_0)} - E[t_0]\dot{x}_{(-\infty, t_0)} + f[t_0] =: \tilde{f}[t_0].$$

The right-hand side can be assumed to be arbitrary, and since  $(E, A)$  is DAE-regular it follows that the operator

$$\begin{aligned} (E \frac{d}{dt} - A)_{t_0} : (\mathbb{D}_{\text{pwc}\infty})^n &\rightarrow \{ D_{t_0} \in (\mathbb{D}_{\text{pwc}\infty})^n \mid \text{supp } D_{t_0} \subseteq \{t_0\} \}, \\ x &\mapsto E_{(t_0, \infty)}\dot{x}[t_0] - A_{(t_0, \infty)}x[t_0] \end{aligned}$$

must be surjective. Assume

$$x[t_0] = \sum_{i=0}^p x_i \delta_{t_0}^{(i)}$$

for some  $p \in \mathbb{N}$  and  $x_0, x_1, \dots, x_p \in \mathbb{R}^n$ , then

$$\dot{x}[t_0] = \sum_{i=0}^{p+1} x_{i-1} \delta_{t_0}^{(i)},$$

where  $x_{-1} := x(t+) - x(t-)$ . Using (10), one gets

$$\begin{aligned}
E_{(t_0, \infty)} \dot{x}[t_0] - A_{(t_0, \infty)} x[t_0] &= \sum_{j=0}^{p+1} \sum_{i=0}^j (-1)^i \binom{j}{i} E^{(i)}(t_0+) x_{j-1} \delta_{t_0}^{(j-i)} \\
&\quad - \sum_{j=0}^p \sum_{i=0}^j (-1)^i \binom{j}{i} A^{(i)}(t_0+) x_j \delta_{t_0}^{(j-i)} \\
&= \sum_{i=0}^{p+1} \sum_{j=i-1}^p (-1)^{j-i+1} \binom{j+1}{i} E^{(j-i+1)}(t_0+) x_j \delta_{t_0}^{(i)} \\
&\quad - \sum_{i=0}^p \sum_{j=i}^p (-1)^{j-i} \binom{j}{i} A^{(j-i)}(t_0+) x_j \delta_{t_0}^{(i)} \\
&= \sum_{i=0}^{p+1} a_i \delta_{t_0}^{(i)} \stackrel{!}{=} \tilde{f}[t_0]
\end{aligned}$$

where

$$\mathbb{R}^{(p+2)n} \ni \begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_{p+1} \end{pmatrix} := \mathcal{N}^{p+1, p+1}(t_0+) \begin{pmatrix} x_{-1} \\ x_0 \\ \vdots \\ x_p \end{pmatrix}.$$

Note that, in particular, for  $i = 0, 1, \dots, p+1$

$$\begin{pmatrix} a_0 \\ a_1 \\ \vdots \\ a_i \end{pmatrix} = \mathcal{N}^{i, p+1}(t_0+) \begin{pmatrix} x_{-1} \\ x_0 \\ \vdots \\ x_p \end{pmatrix}.$$

Since  $a_0, a_1, \dots$ , are given by  $\tilde{f}[t_0]$  they can be arbitrary. Hence there must exist  $q_0 \in \mathbb{N}$  such that  $\mathcal{N}^{0, q_0}$  has full row rank, otherwise not all values for  $a_0$  can be ‘‘produced’’. In general, for every  $i \in \mathbb{N}$  there must exist  $q_i \in \mathbb{N}$  such that  $\mathcal{N}^{i, q_i}$  has full row rank to guarantee that every vector  $(a_0^\top, a_1^\top, \dots, a_i^\top)^\top$  can be obtained. This proves the theorem.

□

## Proof of Theorem 21

- (i) If  $t_0 \geq t_1$ , then the ITP for  $(E, A)$  is identical to the ITP for  $(E_1, A_1)$ , hence only  $t_0 < t_1$  needs to be considered. For  $x^0 \in (\mathbb{D}_{\text{pwc}\infty})^n$  and  $f \in (\mathbb{D}_{\text{pwc}\infty})^n$  let  $x^1$  be the unique solution of the ITP  $(E_0, A_0)$ ,  $x_{(-\infty, t_0)}^1 = x_{(-\infty, t_0)}^0$  with inhomogeneity  $f$  and let  $x$  be the unique solution of the ITP  $(E_1, A_1)$ ,  $x_{(-\infty, t_1)} = x_{(-\infty, t_1)}^1$  with inhomogeneity  $f$ . It will be shown that  $x$  is also the unique solution of the ITP  $(E, A)$ ,  $x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$ . First observe that  $x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^1 = x_{(-\infty, t_0)}^0$  because  $t_0 < t_1$ . Secondly, the following equivalences hold (using Proposition 11)

$$\begin{aligned}
(E\dot{x})_{[t_0, \infty)} &= (Ax + f)_{[t_0, \infty)} \\
\Leftrightarrow (E\dot{x})_{[t_0, t_1)} &= (Ax + f)_{[t_0, t_1)} \quad \wedge \quad (E\dot{x})_{[t_1, \infty)} = (Ax + f)_{[t_1, \infty)} \\
\Leftrightarrow (E_0\dot{x}^1)_{[t_0, t_1)} &= (A_0x^1 + f)_{[t_0, t_1)} \quad \wedge \quad (E_1\dot{x})_{[t_1, \infty)} = (A_1x + f)_{[t_1, \infty)}.
\end{aligned}$$

The last expression is true by the definition of  $x^1$  and  $x$ , hence  $x$  is a solution of the ITP.

It remains to show that  $x$  is unique. Assume there is another solution  $\tilde{x}$ . Since, by definition,  $\tilde{x}_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0 = x_{(-\infty, t_0)}$ , it remains to show that  $\tilde{x}_{[t_0, t_1]} = x_{[t_0, t_1]}$  and  $\tilde{x}_{[t_1, \infty)} = x_{[t_1, \infty)}$ . Let  $z$  and  $\tilde{z}$  be the solutions of the ITP  $(E_0, A_0)$ ,  $z_{(-\infty, t_1)} = x_{(-\infty, t_1)}$  and  $\tilde{z}_{(-\infty, t_1)} = \tilde{x}_{(-\infty, t_1)}$ , resp., then

$$(E_0 \dot{z})_{[t_0, t_1]} = (E_0 \dot{x})_{[t_0, t_1]} = (A_0 x + f)_{[t_0, t_1]} = (A_0 z + f)_{[t_0, t_1]}$$

and

$$(E_0 \dot{\tilde{z}})_{[t_0, t_1]} = (E_0 \dot{\tilde{x}})_{[t_0, t_1]} = (A_0 \tilde{x} + f)_{[t_0, t_1]} = (A_0 \tilde{z} + f)_{[t_0, t_1]}.$$

Hence  $z$  and  $\tilde{z}$  are also solutions of the ITP  $(E_0, A_0)$ ,  $z_{(-\infty, t_0)} = x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$  and  $\tilde{z}_{(-\infty, t_0)} = \tilde{x}_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$ , resp. Since  $(E_0, A_0)$  is DAE-regular it follows that  $z = \tilde{z}$  and therefore  $x_{[t_0, t_1]} = \tilde{x}_{[t_0, t_1]}$ . Finally, observe that  $x$  and  $\tilde{x}$  are solutions of the ITP  $(E_1, A_1)$ ,  $x_{(-\infty, t_1)} = x_{(-\infty, t_1)}$  and  $\tilde{x}_{(-\infty, t_1)} = \tilde{x}_{(-\infty, t_1)}$ , resp. Since  $(E_1, A_1)$  is DAE-regular and  $x_{(-\infty, t_1)} = \tilde{x}_{(-\infty, t_1)}$ , it follows that  $x = \tilde{x}$ .

- (ii) Consider the ITP  $(E, A)$ ,  $x_{(-\infty, \tau_0)} = \xi_{(-\infty, \tau_0)}^0$  for some initial trajectory  $\xi^0$  and  $\tau_0 \in \mathbb{R}$ . Without restriction of generality it may be assumed that  $t_0 \leq \tau_0 < t_1$  (just by changing the indices). Let  $x^0$  be the solution of the ITP  $(E_0, A_0)$ ,  $x_{(-\infty, \tau_0)}^0 = \xi_{(-\infty, \tau_0)}^0$  and, for  $i \in \mathbb{N}$ , let  $x^{i+1}$  be the solution of the ITP  $(E_{i+1}, A_{i+1})$ ,  $x_{(-\infty, t_{i+1})}^{i+1} = x_{(-\infty, t_{i+1})}^i$ . Then  $x = \lim_{i \rightarrow \infty} x^i$  is a well defined distribution and it follows by inductively repeating the same arguments as in (i) that  $x$  is the unique solution of the ITP  $(E, A)$ ,  $x_{(-\infty, \tau_0)} = \xi_{(-\infty, \tau_0)}^0$ . Hence  $(E, A)$  is DAE-regular.
- (iii) Consider the ITP  $(E_0 + E_1[t], A_0 + A_1[t])$ ,  $x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$  for some  $x^0 \in (\mathbb{D}_{\text{pwc}\infty})^n$ ,  $t_0 \in \mathbb{R}$  and with an inhomogeneity  $f \in (\mathbb{D}_{\text{pwc}\infty})^n$ . Clearly, if  $t_0 > t$  this ITP is identical to the ITP  $(E_0, A_0)$  with the same initial trajectory and inhomogeneity. Hence it remains to consider  $t_0 \leq t$ . Let  $\hat{x}$  be the solution of the ITP  $(E_0, A_0)$ ,  $\hat{x}_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$  with inhomogeneity  $f$  and let  $x$  be the solution of the ITP  $(E_0, A_0)$ ,  $x_{(-\infty, t)} = \hat{x}_{(-\infty, t)}$  with inhomogeneity  $\hat{f} := f + A_1[t]\hat{x} - E_1[t]\dot{\hat{x}}$ . It will be shown that  $x$  is the unique solution of the ITP  $(E, A)$ ,  $x_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$  with inhomogeneity  $f$ . First observe that  $x_{(-\infty, t_0)} = \hat{x}_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$ . Secondly,

$$\begin{aligned} ((E_0 + E_1[t])\dot{x})_{[t_0, t]} &= (E_0 \dot{x})_{[t_0, t]} = (E_0 \dot{\hat{x}})_{[t_0, t]} = (A_0 \hat{x} + f)_{[t_0, t]} \\ &= ((A_0 + A_1[t])x + f)_{[t_0, t]} \end{aligned}$$

and, because  $x_{(-\infty, t)} = \hat{x}_{(-\infty, t)}$ ,

$$\begin{aligned} ((E_0 + E_1[t])\dot{x})_{[t, \infty)} &= (E_0 \dot{x})_{[t, \infty)} + E_1[t]\dot{x} \\ &= (A_0 x + \hat{f})_{[t, \infty)} + E_1[t]\dot{\hat{x}} \\ &= (A_0 x + f + A_1[t]\hat{x})_{[t, \infty)} \\ &= ((A_0 + A_1[t])x + f)_{[t, \infty)}. \end{aligned}$$

Hence it remains to show uniqueness of the solution  $x$ . Therefore, let  $\tilde{x}$  also be a solution of the ITP  $(E, A)$ ,  $\tilde{x}_{(-\infty, t_0)} = x_{(-\infty, t_0)}^0$  with inhomogeneity  $f$ . With the same arguments as above it follows that  $\tilde{x}_{(-\infty, t)} = x_{(-\infty, t)} = \hat{x}_{(-\infty, t)}$ . Now

$$\begin{aligned} ((E_0 + E_1[t])\dot{x})_{[t, \infty)} &= ((A_0 + A_1[t])x + f)_{[t, \infty)} \\ \Leftrightarrow (E_0 \dot{x})_{[t, \infty)} &= (A_0 x + \hat{f})_{[t, \infty)} \end{aligned}$$

and the same for  $\tilde{x}$ , hence  $x$  and  $\tilde{x}$  are both solutions of the ITP  $(E_0, A_0)$ ,  $x_{(-\infty, t)} = \hat{x}_{(-\infty, t)}$  with inhomogeneity  $\hat{f}$ . Because  $(E_0, A_0)$  is DAE-regular it follows that  $x = \tilde{x}$ .

- (iv) Let  $T = \{ t_i \in \mathbb{R} \mid i \in \mathbb{Z} \}$  be a locally finite set such that  $E_1[\cdot] = \sum_{i \in \mathbb{Z}} E_1[t_i]$  and  $A_1[\cdot] = \sum_{i \in \mathbb{Z}} A_1[t_i]$ . Furthermore, let  $\tilde{E}_0 = E_0$ ,  $\tilde{A}_0 = A_0$  and, for  $k \in \mathbb{N}$ ,  $\tilde{E}_{k+1} = \tilde{E}_k + E_1[t_k]$ ,  $\tilde{E}_{-k-1} = \tilde{E}_{-k} + E_1[t_{-k}]$ ,  $\tilde{A}_{k+1} = \tilde{A}_k + A_1[t_k]$ ,  $\tilde{A}_{-k-1} = \tilde{A}_{-k} + A_1[t_{-k}]$ . Then it follows inductively from (iii) that  $(\tilde{E}_i, \tilde{A}_i)$  is DAE-regular for all  $i \in \mathbb{Z}$ . Finally,

$$(E_0 + E_1[\cdot], A_0 + A_1[\cdot]) = \left( \sum_{i \in \mathbb{Z}} \tilde{E}_i, \sum_{i \in \mathbb{Z}} \tilde{A}_i \right)$$

and regularity follows from (ii).

□

## Proof of Theorem 24

By Theorem 21 it suffices to consider the impulse free case, i.e.  $E[\cdot] = 0$  and  $A[\cdot] = 0$ .

*The distributional ODE case, i.e.  $(E, A) = (I, A)$ .*

Let  $A^{\text{reg}} \in (\mathcal{C}_{\text{pw}}^\infty)^{n \times n}$  such that  $A_{\mathbb{D}}^{\text{reg}} = A_{\text{reg}} = A$  and consider the standard homogeneous ODE  $\dot{x} = A^{\text{reg}}x$ . Let  $\phi(\cdot, t_0) : \mathbb{R} \rightarrow \mathbb{R}^{n \times n}$ ,  $t_0 \in \mathbb{R}$ , be its fundamental solution, i.e.  $\phi(\cdot, t_0)$  is absolutely continuous,  $\phi(\cdot, t_0)' = A^{\text{reg}}\phi(\cdot, t_0)$  and  $\phi(t_0, t_0) = I$  [27, C.4].

It will be shown first, that  $\phi(\cdot, t_0)$  is piecewise-smooth as in Definition 1. Let  $T = \{ \tau_i \in \mathbb{R} \mid i \in \mathbb{Z} \}$  be a locally finite set such that  $A^{\text{reg}} = \sum_{i \in \mathbb{Z}} A_i$  for some family of smooth matrices  $(A_i)_{i \in \mathbb{Z}}$ . For  $t \in \mathbb{R}$  and  $i \in \mathbb{Z}$  let  $\phi_i(\cdot, t)$  be the fundamental solutions of  $\dot{x} = A_i x$ . Then  $\phi_i(\cdot, t)$  is smooth for all  $i \in \mathbb{Z}$  because each  $A_i$  is smooth. Since the ODEs  $\dot{x} = A^{\text{reg}}x$  and  $\dot{x} = A_i x$  are identical on the interval  $[\tau_i, \tau_{i+1})$  the fundamental solution restricted to this interval are also identical if the initial time fulfills  $t \in [\tau_i, \tau_{i+1})$ , hence  $\phi(s, t) = \phi_i(s, t)$  for all  $s, t \in [t_i, t_{i+1})$ . For a fixed  $t_0 \in \mathbb{R}$  this yields  $\phi(t, t_0) = \phi_i(t, t_i)\phi(t_i, t_0)$  where  $i \in \mathbb{Z}$  is chosen such that  $t \in [t_i, t_{i+1})$ . Now it follows that

$$\phi(\cdot, t_0) = \sum_{i \in \mathbb{Z}} (\phi_i(\cdot, t_i)\phi(t_i, t_0))_{[t_i, t_{i+1})},$$

which shows that  $\phi(\cdot, t_0)$  is piecewise-smooth. Note furthermore, that each  $\phi_i(\cdot, t_i)$  is invertible with  $\phi_i(\cdot, t_i)^{-1} \in (\mathcal{C}^\infty)^{n \times n}$ , this shows that the inverse  $\phi(\cdot, t_0)^{-1}$  is also piecewise-smooth.

It will be shown now that the ITP

$$\begin{aligned} x_{(-\infty, t_0)} &= x_{(-\infty, t_0)}^0, \\ \dot{x}_{[t_0, \infty)} &= (Ax + f)_{[t_0, \infty)}, \end{aligned}$$

where  $x^0, f \in (\mathbb{D}_{\text{pwC}}^\infty)^n$  and  $t_0 \in \mathbb{R}$ , has the unique solution

$$x = x_{(-\infty, t_0)}^0 + \left( \phi(\cdot, t_0)_{\mathbb{D}} x^0(t_0-) + \phi(\cdot, t_0)_{\mathbb{D}} \int_{t_0}^{\cdot} \phi(\cdot, t_0)_{\mathbb{D}}^{-1} f \right)_{[t_0, \infty)}.$$

It must first be shown, that  $\dot{x}_{[t_0, \infty)} = (Ax)_{[t_0, \infty)}$ . Proposition 11, Proposition 12 and  $\phi(\cdot, t_0)'_{\mathbb{D}} = A\phi(\cdot, t_0)_{\mathbb{D}}$  yield

$$\begin{aligned} \dot{x}_{[t_0, \infty)} &= -x^0(t_0-)\delta_{t_0} + \left( \phi(\cdot, t_0)'_{\mathbb{D}} x^0(t_0-) + \left( \phi(\cdot, t_0)_{\mathbb{D}} \int_{t_0}^{\cdot} \phi(\cdot, t_0)_{\mathbb{D}}^{-1} f \right)' \right)_{[t_0, \infty)} \\ &\quad + \left( \underbrace{\phi(\cdot, t_0)_{\mathbb{D}}(t_0-)}_{=I} x^0(t_0-) + \underbrace{\left( \phi(\cdot, t_0)_{\mathbb{D}} \int_{t_0}^{\cdot} \phi(\cdot, t_0)_{\mathbb{D}}^{-1} f \right)}_{=0} (t_0-) \right) \delta_{t_0} \\ &= \left( A\phi(\cdot, t_0)_{\mathbb{D}} x^0(t_0-) + A\phi(\cdot, t_0)_{\mathbb{D}} \int_{t_0}^{\cdot} \phi(\cdot, t_0)_{\mathbb{D}}^{-1} f + f \right)_{[t_0, \infty)} \end{aligned}$$

and, since  $A[t_0] = 0$ ,

$$(Ax + f)_{[t_0, \infty)} = \underbrace{(Ax^0_{(-\infty, t_0)})}_{=0}_{[t_0, \infty)} + \left( A\phi(\cdot, t_0)_{\mathbb{D}} x^0(t_0-) + A\phi(\cdot, t_0)_{\mathbb{D}} \int_{t_0} \phi(\cdot, t_0)_{\mathbb{D}}^{-1} f + f \right)_{[t_0, \infty)},$$

which shows that  $x$  is a solution of the ITP.

It remains to show that the proposed solution is unique. Assume that  $x_1, x_2 \in (\mathbb{D}_{\text{pwc}\infty})^n$  are solutions of the same ITP, then  $e = x_1 - x_2$  fulfills  $e_{(-\infty, t_0)} = 0$  and

$$\dot{e}_{[t_0, \infty)} = (Ae)_{[t_0, \infty)}.$$

Note that  $\dot{e}_{(-\infty, t_0)} = 0 = (Ae)_{(-\infty, t_0)}$ , hence  $e$  is a solution of  $\dot{e} = Ae$  with  $e(t_0-) = 0$  and it must be shown that  $e = 0$  is the only solution of  $\dot{e} = Ae$  with  $e(t_0-) = 0$ . Let  $e \in (\mathbb{D}_{\text{pwc}\infty})^n$  be any solution of  $\dot{e} = Ae$  with  $e(t_0-) = 0$  and let  $\eta := \phi(\cdot, t_0)_{\mathbb{D}}^{-1} e$ . Then, since  $e = \phi(\cdot, t_0)_{\mathbb{D}} \eta$ ,

$$\dot{e} = \underbrace{A\phi(\cdot, t_0)_{\mathbb{D}} \eta}_{=Ae=\dot{e}} + \phi(\cdot, t_0)_{\mathbb{D}} \dot{\eta},$$

hence

$$\dot{\eta} = 0.$$

This implies that  $\eta$ , as an distributional antiderivative of zero, is a constant distribution and since  $\eta(t_0-) = e(t_0-) = 0$  it follows that  $\eta = 0$ . This shows that  $e = 0$  and it is shown that  $x$  as given above is the only solution of the ITP.

*The pure distributional DAE case, i.e.  $(E, A) = (N, I)$  with  $N_{\text{reg}}$  a strictly lower triangular matrix.*

First observe that  $(N\dot{x})_{[t_0, \infty)} = N_{[t_0, \infty)} \dot{x}$ , so the ITP

$$\begin{aligned} x_{(-\infty, t_0)} &= x^0_{(-\infty, t_0)}, \\ (N\dot{x})_{[t_0, \infty)} &= (x + f)_{[t_0, \infty)}, \end{aligned}$$

can equivalently reformulated into a distributional DAE without explicit initial condition:

$$N_{\text{itp}} \dot{x} = x + f_{\text{itp}},$$

where  $N_{\text{itp}} = N_{[t_0, \infty)}$  and  $f_{\text{itp}} = -x^0_{(-\infty, t_0)} + f_{[t_0, \infty)}$ . The matrix  $N_{\text{itp}}$  is still a strictly lower triangular matrix. Consider the operator

$$N_{\text{itp}} \frac{d}{dt} : (\mathbb{D}_{\text{pwc}\infty})^n \rightarrow (\mathbb{D}_{\text{pwc}\infty})^n, \quad x \mapsto N_{\text{itp}} \dot{x},$$

and its powers

$$(N_{\text{itp}} \frac{d}{dt})^0 := x \mapsto x, \quad \forall i \in \mathbb{N} : (N_{\text{itp}} \frac{d}{dt})^{i+1} := x \mapsto (N_{\text{itp}} \frac{d}{dt}) \left( (N_{\text{itp}} \frac{d}{dt})^i(x) \right).$$

Then, since  $N_{\text{itp}}$  is a strictly lower triangular matrix, the operator  $N_{\text{itp}} \frac{d}{dt}$  is nilpotent, i.e. there exists  $\nu \in \mathbb{N}$  such that  $(N_{\text{itp}} \frac{d}{dt})^\nu$  is the zero operator. From this it follows that the operator

$$(N_{\text{itp}} \frac{d}{dt} - I) : (\mathbb{D}_{\text{pwc}\infty})^n \rightarrow (\mathbb{D}_{\text{pwc}\infty})^n, \quad x \mapsto N_{\text{itp}} \dot{x} - x$$

is bijective with inverse

$$(N_{\text{itp}} \frac{d}{dt} - I)^{-1} = - \sum_{i=0}^{\nu-1} (N_{\text{itp}} \frac{d}{dt})^i.$$

Hence the unique solution of the ITP is given by

$$x = - \sum_{i=0}^{\nu-1} \left( N_{[t_0, \infty)} \frac{d}{dt} \right)^i \left( -x^0_{(-\infty, t_0)} + f_{[t_0, \infty)} \right).$$

□

---

## References

1. Campbell, S.L.: Singular Systems of Differential Equations II. Pitman, New York (1982)
2. Cobb, J.D.: Controllability, observability and duality in singular systems. *IEEE Trans. Autom. Control* **AC-29**, 1076–1082 (1984)
3. Colombeau, J.F.: Multiplication of distributions. No. 1532 in *Lecture Notes in Mathematics*. Springer Verlag, Berlin Heidelberg (1992)
4. Doležal, V.: The existence of a continuous basis of a certain linear subspace of  $E_r$ , which depends on a parameter. *Časopis pro pestovani matematiky* **89**, 466–468 (1964)
5. Fisher, B., Lin-Zhi, C.: The product of distributions on  $R^n$ . *Commentat. Math. Univ. Carol.* **33**(4), 605–614 (1992)
6. Fuchssteiner, B.: Eine assoziative Algebra über einen Unterraum der Distributionen. *Mathematische Annalen* **178**, 302–314 (1968)
7. Fuchssteiner, B.: Algebraic foundation of some distribution algebras. *Studia Mathematica* **76**, 439–453 (1984)
8. Geerts, A.H.W.T.: Solvability conditions, consistency and weak consistency for linear differential-algebraic equations and time-invariant linear systems: The general case. *Lin. Alg. Appl.* **181**, 111–130 (1993)
9. Geerts, A.H.W.T.: Higher-order continuous-time implicit systems: Consistency and weak consistency, impulse controllability, geometric concepts, and invertibility properties. *Lin. Alg. Appl.* **244**, 203–253 (1996)
10. Geerts, A.H.W.T., Schumacher, J.M.: Impulsive-smooth behavior in multimode systems. Part I: State-space and polynomial representations. *Automatica* **32**(5), 747–758 (1996)
11. Geerts, A.H.W.T., Schumacher, J.M.: Impulsive-smooth behavior in multimode systems. Part II: Minimality and equivalence. *Automatica* **32**(6), 819–832 (1996)
12. Hautus, M.L.J.: The formal Laplace transform for smooth linear systems. In: G. Marchesini, S.K. Mitter (eds.) *Mathematical Systems Theory*, no. 131 in *Lecture Notes in Economics and Mathematical Systems*, pp. 29–47. Springer-Verlag, New York (1976)
13. Hautus, M.L.J., Silverman, L.M.: System structure and singular control. *Lin. Alg. Appl.* **50**, 369–402 (1983)
14. Hinrichsen, D., Prätzel-Wolters, D.: Solution modules and system equivalence. *Int. J. Control* **32**, 777–802 (1980)
15. Hoskins, R.F., Sousa Pinto, J.J.M.: Distributions, ultradistributions and other generalized functions. *Mathematics and its applications*. Ellis Horwood, Chichester, West Sussex, UK (1994)
16. Hou, M., Pugh, A.C., Hayton, G.E.: A test for behavioral equivalence. *IEEE Trans. Autom. Control* **45**(11), 2177–2182 (2000)
17. König, H.: Multiplikation von Distributionen. I. *Mathematische Annalen* **128**, 420–452 (1955)
18. König, H., Raeder, R.: Vorlesung über die Theorie der Distributionen. No. 6-1 in *Annales Universitatis Saraviensis - Series Mathematicae*. Universität des Saarlandes, Saarbrücken (1995)
19. Kunkel, P., Mehrmann, V.: *Differential-Algebraic Equations. Analysis and Numerical Solution*. EMS Publishing House, Zürich, Switzerland (2006)
20. Lakshmikantham, V., Bainov, D.D., Simeonov, P.S.: *Theory of Impulsive Differential Equations*. No. 6 in *Series in Modern Applied Mathematics*. World Scientific, Singapore, Teaneck, NJ (1989)
21. Liberzon, D.: *Switching in Systems and Control. Systems and Control: Foundations and Applications*. Birkhäuser, Boston (2003)
22. Mikusiński, J.: On the square of the Dirac delta-distribution. *Bull. Acad. Pol. Sci., Sér. sci. math. astr. phys.* **XIV**(9), 511–513 (1966)
23. Oberguggenberger, M.: Multiplication of distributions and applications to partial differential equations. No. 259 in *Pitman research notes in mathematics*. Longman Scientific & Technical, Harlow, Essex, UK (1992)
24. Rabier, P.J., Rheinboldt, W.C.: Time-dependent linear DAEs with discontinuous inputs. *Lin. Alg. Appl.* **247**, 1–29 (1996)

- 
25. Rabier, P.J., Rheinboldt, W.C.: Theoretical and numerical analysis of differential-algebraic equations. In: P.G. Ciarlet, J.L. Lions (eds.) *Handbook of Numerical Analysis*, vol. VIII, pp. 183–537. Elsevier Science, Amsterdam, The Netherlands (2002)
  26. Schwartz, L.: *Théorie des Distributions*. Hermann, Paris (1957, 1959)
  27. Sontag, E.D.: *Mathematical Control Theory, Deterministic Finite Dimensional Systems*, 2nd edn. Springer-Verlag, New York (1998)
  28. Vardulakis, A.I.G., Antoniou, E.N., Karampetakis, N.: On the solution and impulsive behaviour of polynomial matrix descriptions of free linear multivariable systems. *Int. J. Control* **72**(3), 215–228 (1999)
  29. Vardulakis, A.I.G., Fragulis, G.: Infinite elementary divisors of polynomial matrices and impulsive solutions of linear homogeneous matrix differential equations. *Circuits Systems Signal Process.* **8**(3), 357–373 (1989)
  30. Verghese, G.C., Levy, B.C., Kailath, T.: A generalized state-space for singular systems. *IEEE Trans. Autom. Control* **AC-26**(4), 811–831 (1981)
  31. Walter, H.F.: Über die Multiplikation von Distributionen in einem Folgenmodell. *Mathematische Annalen* **189**, 211–221 (1970)
  32. Walter, W.: *Einführung in die Theorie der Distributionen*, 3rd edn. BI Wissenschaftsverlag, Mannheim Leipzig Wien Zürich (1994)
  33. Weierstraß, K.: Zur Theorie der bilinearen und quadratischen Formen. *Monatsh. Akademie. Wiss.* pp. 310–338 (1868)
  34. Willems, J.C.: The behavioral approach to open and interconnected systems. *IEEE Control Systems Magazine* **27**(6), 46–99 (2007)
  35. Yip, E.L., Sincovec, R.F.: Solvability, controllability and observability of continuous descriptor systems. *IEEE Trans. Autom. Control* **AC-26**, 702–707 (1981)